arXiv:1005.0321v2 [quant-ph] 4 Jul 2011

Quantum theory for a total system with one internal measuring apparatus

Wen-ge Wang
Department of Modern Physics, University of Science and Technology of China, Hefei 230026, China

(Dated: July 5, 2011)

In this paper, we generalize the standard formalism of quantum mechanics to a quantum theory for a total system including one internal measuring apparatus. The generalization is mainly based on the limitedness of the measuring ability of the internal measuring apparatus, that is, the internal measuring apparatus can not measure all its own properties; this limitedness implies the possibility that measurements performed by the internal measuring apparatus can not distinguish between some pure-vector descriptions and some mixed-state descriptions of the total system. The proposed theory has three basic assumptions, which roughly speaking have the following contents: (i) Physical states of the total system can be associated with vectors in the total Hilbert space; (ii) the dynamical evolution of a state vector obeys Schrödinger equation; and (iii) under a principle of compatible description and certain non-transition condition, a pure-vector description of the total system may imply the existence of certain mixed-state description. The principle of compatible description states that different mathematical descriptions for the same physical state of the total system must give consistent predictions for results of measurements performed by the internal measuring apparatus. This principle imposes a restriction to vectors in the Hilbert space and this may effectively break the time-reversal symmetry of Schrödinger equation. As applications of the proposed theory, the possibility for an n-level system to possess definite properties is discussed and a master equation is derived for the time evolution of a system that may have definite properties for most of the times.

PACS numbers: 03.65.Ta; 03.65.-w; 03.65.Yz

Contents

I. Introduction 2

II. Preliminary analysis and structure of the paper 3
  A. An implication of the limited measuring ability of the internal measuring apparatus 3
  B. Basic contents and structure of the paper 3

III. Basic assumptions of the theory 4
  A. The first and second basic assumptions 4
  B. General form of the third assumption 5

IV. Explicit form of Part II of the third assumption 5
  A. A technique to be used 5
  B. Definite properties described in the Hilbert space — R-observable 6
    1. Formal expression of definite property 6
    2. R-observable 6
    3. Fine and coarse-grained R-observables 7
  C. Explicit form of the condition C_D 7
  D. Assumption of mixed-state description 8

V. A concise expression of the third assumption 8
  A. Tree structure of branching for time evolution 8
  B. Fine and coarse-grained trees 10
  C. A requirement of the principle of compatible description 11
  D. Initial-vector-restriction and a concise expression of the third assumption 12
  E. Further discussions in the condition C_D and the possibility of experimental test 13

VI. Measuring apparatus and measurement 13

VII. Decoherence Mechanism 14
  A. Time evolution under the non-transition condition 14
  B. Decoherence mechanism for the existence of R-observable 14
  C. $\mathcal{D}_{\alpha\alpha'}$ expressed as a generalized Loschmidt echo 15

VIII. Application I — Isolatable and non-isolatable systems 16

IX. Application II — An n-level system 17
  A. Energy eigenstates and R-observable 17
  B. $D_{\beta\alpha}$ expressed in the Hilbert space of the environment 18

X. Application III — Irreversibility of branching processes 18
  A. Irreversible feature of time evolution 18
  B. Master equation for an ideal case of branching 18

XI. Discussions and conclusions 19
  A. Comparison with CHI of quantum mechanics 19
  B. Comparison with MWI of quantum mechanics 20
  C. Summary and discussions 21

Acknowledgments 21

A. A relation between trees and their coarse-grainings 21

B. R-observable and preferred pointer basis 22

References 22

I. INTRODUCTION

The standard formalism of quantum mechanics [1], which has passed all experimental tests ever performed, is basically a theory for an external observer; it gives predictions for measurements performed by an external measuring apparatus. A topic that has received extensive attention since the establishment of the formalism, with lots of controversy, is the possibility of extending it to a quantum theory for an isolated, total system (like the universe) described by an internal observer. To achieve this goal, the major difficulty comes from treatment of the measuring apparatus, which is now a part of the total system. This difficulty is related to the so-called measurement problem, concerning the relationship between Schrödinger evolution and definite outcomes of measurements.

The above mentioned problem is of interest not only for pure theoretical reasons, but also for a practical reason, concerning designation of small measuring apparatus. In recent years, significant progresses have been achieved in technology, such that it is now a commonplace in labs to observe small systems at the mesoscopic scale, even at the microscopic scale. In principle, it is possible to design a measuring apparatus, whose essential part is of the microscopic scale. A challenging task is to know the condition under which a small quantum system may possess some definite properties. Our intuition obtained in the macroscopic world is not so helpful for this purpose.

Lots of efforts have been seen in the attempt of solving the above mentioned problem (see, e.g., reviews given in Refs. [2–4]), most under the name of interpretation of quantum mechanics, for example, various versions of Everett's relative-state interpretation (RSI) [5–7], consistent-histories interpretations (CHI) [8–16]) first proposed by Griffiths [8], and De Broglie's pilot wave theory [17] and Bohmian mechanics [18, 19].

There existing so many theories, what is the reason for us to intend to develop another one? Besides the fact that there is no commonly-accepted solution to the measurement problem yet, a major motivation is as follows. Since Schrödinger equation has passed all experimental tests ever performed, here we take the relatively conservative viewpoint that this equation and the corresponding Hilbert space can be taken as a starting point. One may note that this starting point is also taken in RSI and effectively so in CHI [16]. As to be discussed below, although both of RSI and CHI supply quite general frameworks for quantum descriptions, neither of them gives a concrete condition, under which a considered system may have a definite property. To find such a concrete condition is the main motivation of our work.

Due to the above-mentioned starting point we are to adopt, in this paper we are not to discuss approaches like the dynamical-reduction models [20–25], which introduce modifications to Schrödinger equation. We are not to discuss the approach of Bohm, either, because in addition to a description given in the Hilbert space, this approach requires a hidden variable, i.e., positions of particles. Here we consider only descriptions given in the Hilbert space.

For RSI, to be specific, let us discuss a special version of it, namely, the many-worlds interpretation (MWI) of quantum mechanics. In the theory of MWI, when the state vector for a whole system splits into branches, each branch may have a property more definite than that of the whole state vector. Thus, in principle, the theory allows the appearance of some definite property of a system. However, no concrete condition has been given yet for branching to happen in MWI.

It has been suggested that a combination of MWI and the decoherence theory [26, 27] (see reviews given in [2, 4, 28]) may do better, with preferred (pointer) states in the decoherence theory related to branches in MWI [28]. However, a concrete condition for branching to happen is still missing, because this approach faces the following problem, which is related to the fact that decoherence in the decoherence theory is indicated by vanishing of off-diagonal elements of the reduced density matrix of a subsystem in certain (preferred) basis. That is, a measuring apparatus may be repeatedly used and, during each measuring process, some off-diagonal elements of its reduced density matrix may become non-negligible due to its interaction with the measured system. Therefore, at least for repeatedly used measuring apparatuses, off-diagonal elements of their reduced density matrices can not vanish forever. Relatedly, the decoherence theory faces a problem stressed by A.J.Leggett, namely, distinguishing 'false' and 'true' decoherence [29].

The theory of CHI considers consistent histories. In this theory, when a measuring process is described by a consistent history within a fixed framework, a subsystem may have certain definite properties at some times. However, starting from a given initial condition, there may exist many incompatible frameworks, giving incompatible descriptions (histories). To avoid inconsistency, a single-framework rule is assumed, which states that one is allowed to adopt only one framework in a consistent discussion, though other frameworks are equally valid. (The physical origin of this single-framework rule is still not clear and there have been some debates in its validity [30, 31].) Therefore, the present form of CHI does not supply a method of selecting a specific framework from the many incompatible frameworks [32], as a result, it does not give a condition under which a definite property of a measuring apparatus may appear. In the next section, we discuss our approach to the problem.

II. PRELIMINARY ANALYSIS AND STRUCTURE OF THE PAPER

As discussed above, we take Schrödinger equation and the Hilbert space as our staring point. As shown in von Neumann's analysis [1], these two assumptions are not sufficient for describing measurement. Therefore, a third assumption is needed, which enables one to describe measurement outcomes [33]. The main task of this paper is to find such a third assumption for a total system including one internal measuring apparatus. In this section, we first give some preliminary analyses in properties of the internal measuring apparatus, then, discuss the strategy to be adopted, as well as the basic structure of the paper.

A remark: For the simplicity in discussion, in this paper, we focus on the case that the total system has one internal observer only [34]. Since measuring apparatuses that can be controlled by one observer may always be regarded as forming a big measuring apparatus, without the loss of generality, we consider the case that there exists only one internal measuring apparatus.

A. An implication of the limited measuring ability of the internal measuring apparatus

The main difference between a quantum theory for a total system including one internal measuring apparatus and the usual quantum mechanics lies in the internality of the measuring apparatus, i.e., in the former case the measuring apparatus is a part of the described system. To understand implications of this point, we need to give a briefly discussion in the most essential feature of a measuring apparatus, that is, it may have some definite properties that may be used as measurement records. Let us use μ to denote a possible, definite property of the measuring apparatus. To describe, within the Hilbert space, the fact that a measuring apparatus has a definite outcome after a measurement, typically, one assumes that the whole system has a mixed-state description after the measurement, which we denote by ρ. For example, in the case that with a probability p_μ the whole system is described by a normalized vector $|\Psi_\mu\rangle$, the mixed-state description is written as $\rho = \sum p_\mu |\Psi_\mu\rangle\langle\Psi_\mu|$.

The main problem one needs to solve in order to develop a quantum theory for a total system including the measuring apparatus, is to find a relation between the above discussed mixed-state description ρ and Schrödinger evolution $|\Psi(t)\rangle$, which is just the essence of the measurement problem. We observe that a clue to a solution to this problem is given by a consequence of the internality of the internal measuring apparatus, namely, the limitedness of its measuring ability.

In fact, an internal measuring apparatus can not measure all properties of the total system, at least, it can not measure all its own properties, therefore, there is an intrinsic limitation in its measuring ability. This suggests the possibility that measurements performed by the internal measuring apparatus may not be able to distinguish between some pure-vector description $|\Psi\rangle$ and some mixed-state description ρ of the total system. Indeed, if there may exist two descriptions, which, as well as their time evolutions, always give consistent predictions for results of measurements performed by the internal measuring apparatus, then, in view of the internal observer, the assumption that the two descriptions can be associated with the same physical state of the total system will not lead to any physical inconsistency. We'll use this assumption as the basic content of the third assumption in the theory to be proposed [35].

B. Basic contents and structure of the paper

The theory to be proposed is based on three basic assumptions. In Sec. III, we give the first two basic assumptions (about the Hilbert space and Schrödinger equation), as well as a general form of the third assumption with contents discussed in the last paragraph of the previous section. In particular, we call the following requirement the principle of compatible description, that is, two descriptions for the same physical state of the total system must give consistent predictions for results of measurements performed by the internal measuring apparatus. Thus, basically, the third assumption states that, under the principle of compatible description as well as some other (to be determined) conditions, there may exist certain relation between some mixed-state descriptions and some pure-vector descriptions for the total system.

A main task of this paper is to find out an explicit form of the third assumption, which will be done in Sec.IV. We'll do this based on the following two observations. (1) Our experiences obtained in labs show that a recordable property of a measuring apparatus should have two main features, namely, definiteness and some type of initial-state independence. (2) It would be consistent to use a part of the standard formalism of quantum mechanics, if this part may turn out to be derivable from the theory to be proposed. We'll show that these two points may be used to determine most of the contents of an explicit form of the third assumption, apart from a concrete condition for a definite property of a considered system to appear; the two points can determine only a necessary part of the condition. Finally, resorting to a decoherence consideration, we'll propose the content of the concrete condition and get an explicit form of the third assumption.

In Sec. V, we derive a mathematical expression for the principle of compatible description. It imposes a restriction to the initial condition, which we call Initial-vector-restriction. Making use of this result, the third assumption can be expressed in a concise form. The above discussed strategy of developing the theory is schematically plotted in Fig. 1. In the case that an initial condition can not pass the consistency-checking indicated in the figure, the considered subsystem can not be employed as


(A) Internality of measuring apparatus implies limited measuring ability
suggest
(B) Assumption for the multiplicity of descriptions (|Ψ>,ρ)
predict
(C) Definite property of a subsystem
imply
(D) Records of measurement
admit
(H) Initial condition
impose restriction to
(F) Physical consistency of (B)
check
(E) Principle of compatible description
define
(G) Measuring apparatus

FIG. 1: Schematic plot for a strategy of solving the measurement problem for a total system including one internal measuring apparatus. It starts from the internality of the measuring apparatus, which implies the limitedness of its measuring ability (A). This opens the possibility for a physical state of the total system to have both a pure-vector description and a mixed-state description at the same time (B). The mixed-state descriptions may be used to predict some definite properties of some subsystem (C), which may give measurement records (D). With the help of the predicted measurement records, one may check the physical consistency of the multi-descriptions given by the assumption in part B (F), by making use of the principle of compatible description (E). In the case that the consistency-checking is passed, the considered subsystem may be used as an essential part of a measuring apparatus (G). This imposes a restriction to the initial condition (H), i.e., the assumption in part B is applicable only for those initial conditions that may pass the consistency-checking.

a measuring apparatus. This is not strange. Indeed, according to our experiences, not for all initial states of a total system, could a subsystem of it be used as a measuring apparatus. For example, a subsystem may not be stable enough to be used as a measuring apparatus, when the temperature is too high.

Thereafter, we use the proposed theory to discuss various topics. In Sec.VI, we give a general discussion for measurement and show that the axiom of measurement in the standard formalism of quantum mechanics can be derived under some appropriate conditions. In Sec.VII, we discuss the role played by decoherence; in particular, decoherence supplies a dynamical mechanism for the principle of compatible description to be satisfied.

Sections VIII, IX, and X are devoted to some applications of the proposed theory. In Sec. VIII, we show that the third assumption is not applicable to an isolatable system; this implies that it is relatively easy for an isolatable system (for example, the center-of-mass degrees of freedom of a system) to keep coherence. In Sec.IX, we discuss an n-level system and show that it may have definite properties under adequate conditions. The last application is given in Sec.X for irreversible features of some processes. The irreversibility comes from the restriction imposed by the principle of compatible description to the initial condition, which may effectively break the time-reversal symmetry of Schrödinger equation for some processes.

Finally, discussions and conclusions are given in Sec.XI. In particular, we discuss the main similarities and differences between the proposed theory and CHI and MWI of quantum mechanics.

III. BASIC ASSUMPTIONS OF THE THEORY

In this paper, we consider an isolated, total system, which is composed of a system $\mathcal{R}$ and its environment denoted by $\mathcal{E}$. It is the system $\mathcal{R}$ that is expected to be able to play the role of the internal measuring apparatus. To avoid confusion, we would stress that, in the logic structure of the theory to be proposed, the concept of measuring apparatus is not a fundamental concept, but, is a defined concept. (The definition will be given in Sec.VI.)

In this section, we introduce the first two basic assumptions and a general form of the third basic assumption of the theory to be proposed. We assume that one does not need to give the internal observer a special position at the fundamental level of the theory. Thus, the internal observer may be regarded as a part of the environment $\mathcal{E}$.

A. The first and second basic assumptions

As mentioned in the section of introduction, we follow the usual quantum mechanics for the first two basic assumptions. The first one is about the state space.

- **Postulate of Hilbert space (HS)**: Each physical state of an isolated system, which is described by an internal observer, can be associated with a vector or a density operator in the total Hilbert space $\mathscr{H}$.

We use $\mathscr{H}_\mathcal{R}$, $\mathscr{H}_\mathcal{E}$, and $\mathscr{H}$ to denote the Hilbert spaces corresponding to the system $\mathcal{R}$, its environment $\mathcal{E}$, and the total system $\mathcal{R}+\mathcal{E}$, respectively, with $\mathscr{H} = \mathscr{H}_\mathcal{R} \otimes \mathscr{H}_\mathcal{E}$. A density operator is given the ordinary ensemble interpretation. We remark that extension of the Hilbert space considered in the usual quantum mechanics, which is for descriptions given by an external observer, to the total Hilbert space discussed in the postulate of HS is a non-trivial extension.

The second basic assumption is about the dynamical law.

- **Postulate of Schrödinger equation (SE)**: The time evolution of a vector description $|\Psi(t)\rangle$ of a physical state of the total system $\mathcal{R}+\mathcal{E}$ obeys Schrödinger equation,

$$\mathrm{i}\hbar\frac{\partial}{\partial t}|\Psi(t)\rangle = H|\Psi(t)\rangle. \tag{1}$$

Here, we use H to denote the Hamiltonian of the total system $\mathcal{R}+\mathcal{E}$,

$$H = H_{\mathcal{R}} + H_{\mathcal{E}} + H_I, \tag{2}$$

where $H_{\mathcal{R}}$ and $H_{\mathcal{E}}$ are the Hamiltonians of $\mathcal{R}$ and $\mathcal{E}$, respectively, and H_I indicates the interaction between $\mathcal{R}$ and $\mathcal{E}$. We use $U(t,t_0)$ to denote the unitary evolution operator,

$$|\Psi(t)\rangle = U(t,t_0)|\Psi(t_0)\rangle. \tag{3}$$

In the case of a time-independent Hamiltonian, $U(t,t_0) = e^{-\mathrm{i}H(t-t_0)/\hbar}$.

B. General form of the third assumption

As discussed in Sec. II, the third assumption is introduced, based on the following fact: The necessary and sufficient condition for a physical state of the total system $\mathcal{R}+\mathcal{E}$ to have two mathematical descriptions at the same time, is that the two descriptions are experimentally compatible with respect to the internal measuring apparatus. Since a measurement record is essentially a definite property of the measuring apparatus, we express this fact in the form of the following principle.

- **The principle of compatible description**: Different mathematical descriptions for the same physical state of the total system $\mathcal{R}+\mathcal{E}$ must give compatible predictions for the probabilities for the system $\mathcal{R}$ to have definite properties.

Obviously, this principle must be obeyed in all physical theories.

Then, summarizing the discussions given in Sec.II A, generally to say, the third assumption may have the following contents: That is, subject to requirements like the principle of compatible description, some pure-vector description $|\Psi\rangle$ and some mixed-state description ρ may describe the same physical state of the total system. More explicitly, its general form can be stated as follows.

General form of the third assumption:

- Part I: The Part II of this assumption is applicable to a vector $|\Psi\rangle$, if the application of Part II to this vector and its time evolutions does not lead to confliction with the principle of compatible description.
- Part II: For a physical state of the total system $\mathcal{R}+\mathcal{E}$ described by a vector $|\Psi\rangle$, when certain condition denoted by C_D is satisfied, the same physical state also has certain mixed-state description.

An explicit form of Part II of the third assumption will be discussed and given in the next section.

Below, we give some further remarks and comments regarding the general form of the third assumption.

(1) Part I of the third assumption guarantees the physical consistency of the theory. That is, only those descriptions that satisfy the principle of compatible description may be associated with the same physical state of the total system.

(2) Composed of two involved parts, the third assumption has an unusual mathematical structure, not possessed by usual physical theories. On one hand, the practical applicability of Part II of the assumption is subject to Part I. On the other hand, satisfaction of Part I of the assumption is determined by whether predictions of Part II obey the principle of compatible description. This feature has its origin in the two-fold roles played by the internal measuring apparatus, namely, it is a part of the described total system, meanwhile it is the system that may check the physical consistency of descriptions of the total system.

(3) In the usual quantum mechanics, it is straightforward to show that $|\psi\rangle$ and $e^{\mathrm{i}\theta}|\psi\rangle$ always satisfy the principle of compatible description, therefore, it is unnecessary to explicitly mention this principle there. However, in the theory here, with the possibility that a same physical state may have a pure-vector description and a mixed-state description at the same time, special attention much be paid to this principle.

IV. EXPLICIT FORM OF PART II OF THE THIRD ASSUMPTION

In this section, we discuss un-specified elements in Part II of the general form of the third assumption given above, namely, the condition C_D and the mixed-state description, and propose an explicit form of Part II of the third assumption. As discussed in Sec.II B, we'll do this, based on a part of the standard formalism of quantum mechanics and two features of measuring apparatuses, namely, definiteness and some type of initial-state independence.

A. A technique to be used

In this section, we discuss a technical method that will be used in our discussion of the explicit form of the third assumption. The method is that, in principle, we may imagine the existence of an external observer, who has no interaction with the composite system $\mathcal{R}+\mathcal{E}$, such that $\mathcal{R}+\mathcal{E}$ is still isolated. Temporarily, we assume that some part of the standard formalism of quantum mechanics can be used by the imaginary external observer to give predictions for the composite system $\mathcal{R}+\mathcal{E}$. For the self-consistency of the results to be obtained, later, we'll show that the temporary assumption can be derived from the theory to be proposed, when the imaginary external observer is regarded as a part of a big system which also includes $\mathcal{R}+\mathcal{E}$ (see Sec.V D).

Specifically, we make the following temporary assumption $\mathrm{A_T}$ for the imaginary external observer.

- $\mathrm{A_T}$. In the case that the internal observer predicts that the system $\mathcal{R}$ may have a definite property, the imaginary external observer may use the axiom of measurement in the standard formalism of quantum mechanics to predict measurement results for the corresponding observable of $\mathcal{R}$.

In principle, the imaginary external observer may communicate with the internal observer for checking the information they have, therefore, the two observers must give compatible predictions for definite properties of the system $\mathcal{R}$.

In addition, note that the following assumption does not conflict with the postulates of HS and of SE introduced in the previous section. That is, the imaginary external observer may use the same pure-vector description $|\Psi(t)\rangle$ as the internal observer to describe the state of $\mathcal{R}+\mathcal{E}$.

B. Definite properties described in the Hilbert space — R-observable

In this section, we discuss mathematical description of definite and recordable properties of a measuring apparatus in the Hilbert space. We first discuss a formal description and make use of it to re-express Part II of the general form of the third assumption. Then, we discuss main features of the definite properties and introduce a concept of R-observable to characterize these features. Finally, we show that each system $\mathcal{R}$ has a finest R-observable.

1. *Formal expression of definite property*

A definite and recordable property of a measuring apparatus $\mathcal{R}$ can be labeled by a quantity with discrete values, which we denote by μ as done above. The simplest way of describing such a property in the Hilbert space is to assume that it corresponds to a division of the Hilbert space of $\mathcal{R}$ into orthogonal subspaces [1], which we denote by $\mathscr{H}_{\mathcal{R}\mu}$. Let us use $\mathcal{P}_\mu$ to indicate the corresponding projection operators for the subspaces, which satisfy

$$\mathcal{P}_\mu \mathcal{P}_\nu = \delta_{\mu\nu}\mathcal{P}_\mu, \qquad \sum_\mu \mathcal{P}_\mu = I_{\mathcal{R}}, \tag{4}$$

where $I_{\mathcal{R}}$ is the identity operator in $\mathscr{H}_{\mathcal{R}}$. Correspondingly, the total Hilbert space is also divided into a series of subspaces

$$\mathscr{H}_\mu \equiv \mathscr{H}_{\mathcal{R}\mu} \otimes \mathscr{H}_{\mathcal{E}}. \tag{5}$$

Obviously, $\mathcal{P}_\mu \otimes I_{\mathcal{E}}$ is the projection operator for the subspace $\mathscr{H}_\mu$, where $I_{\mathcal{E}}$ is the identity operator in $\mathscr{H}_{\mathcal{E}}$. For brevity, without the risk of confusion, we also use $\mathcal{P}_\mu$ to indicate $\mathcal{P}_\mu \otimes I_{\mathcal{E}}$ in what follows. Corresponding to a set of projection operators $\mathcal{P}_\mu$, one may introduce an observable $A_{\{\mu\}}$ for the system $\mathcal{R}$,

$$A_{\{\mu\}} = \sum_\mu \mu \mathcal{P}_\mu. \tag{6}$$

Now, we re-express Part II of the general form of the third assumption. For a reason that will become clear later, we assume that the condition C_D in the general form of the third assumption is satisfied for a time period, not for an instant of time. Without the loss of generality, we use $\mathcal{T}=[0,T]$ to indicate this time period. Then, the Part II may be expressed in the following way.

- Part II: If a condition C_D is satisfied for a description $|\Psi(t)\rangle$ of a physical state of the total system $\mathcal{R}+\mathcal{E}$ for times $t\in\mathcal{T}$, then, at the time $t=T$ the physical state of the total system also has the following mixed-state description, namely, with certain probability p_μ the total system is described by certain vector $|\Psi_\mu\rangle \in \mathscr{H}_\mu$, written as,

$$\rho(T) = \sum_\mu p_\mu |\Psi_\mu\rangle\langle|\Psi_\mu|. \tag{7}$$

2. *R-observable*

In this section, we discuss properties of the projection operators $\mathcal{P}_\mu$, which may correspond to definite properties of a measuring apparatus. We term the corresponding observable $A_{\{\mu\}}$ a *R-observable* of the system $\mathcal{R}$. According to our experiences obtained in labs (also in daily-life), measurement records of a measuring apparatus have the following main features:

- Definiteness.
- Certain type of initial-state independence. (For certain type of initial condition, a measuring apparatus may have the same type of definite property, corresponding to the same set of projection operators $\mathcal{P}_\mu$.)

These features imply that not an arbitrary Hermitian operator can be regarded as a R-observable. Below, we analyze these two features and propose an explicit definition for the concept of R-observable.

First, we discuss the requirement of definiteness, with the help of the method discussed in Sec. IV A. In view of the imaginary external observer, all measurable properties of the system $\mathcal{R}$ can be given by the reduced density matrix $\rho^{re}_{\mathcal{R}}(t) = \mathrm{Tr}_{\mathcal{E}}|\Psi(t)\rangle\langle\Psi(t)|$. Suppose, in view of the internal observer, the system $\mathcal{R}$ has a definite value of μ with certain probability within some time period. Then, for the consistency of the descriptions given by the two observers, $\rho^{re}_{\mathcal{R}}(t)$ must be block-diagonal with respect to the subspaces $\mathscr{H}_{\mathcal{R}\mu}$, namely, $\mathcal{P}_\mu \rho^{re}_{\mathcal{R}}(t) \mathcal{P}_\nu \propto \delta_{\mu\nu}$ for the same time period. This property of the reduced density matrix is usually referred to as decoherence induced by environment.

More exactly, the consistency of the two descriptions discussed above requires that $\mathcal{P}_\mu \rho^{re}_{\mathcal{R}}(t) \mathcal{P}_\nu \doteq 0$ for $\mu \neq \nu$. Here, $A \doteq B$ means $|A - B| \le \epsilon_x$ (or $\|A - B\| \le \epsilon_x$, or the like), where $\epsilon_x \ge 0$ is a small quantity such that its difference from zero generates no effect that may be tested by experiments. Thus, from the viewpoint of experimental test, $A \doteq B$ is effectively equivalent to $A = B$ [36].

Next, we discuss the requirement of initial-state independence. We should be careful here, since due to the time-reversal symmetry of Schrödinger equation one can not expect that decoherence may happen within a finite time period for all initial conditions. In fact, each vector $|\Psi\rangle$ in the Hilbert space has a time-reversal vector, which we denote by $|\widetilde{\Psi}\rangle$; if off-diagonal elements of the reduced density matrix of the system $\mathcal{R}$ decrease when the state vector evolves from $|\Psi_0\rangle$ into $|\Phi\rangle$ within a time period T, it may happen that off-diagonal elements of the reduced density matrix increase within a time period T when $|\widetilde{\Phi}\rangle$ is taken as the initial vector.

Therefore, we can not require an absolute initial-vector independence of the R-observable. Since increasing of off-diagonal elements of the reduced density matrix is usually due to some coherence possessed by the initial vector, in order to determine R-observable, we may consider initial uncorrelated states, which are described by direct products of vectors in the Hilbert spaces of $\mathcal{R}$ and $\mathcal{E}$, respectively, namely, $|\Psi(0)\rangle = |\psi^{\mathcal{R}}_0\rangle |\phi^{\mathcal{E}}_0\rangle$ with $|\psi^{\mathcal{R}}_0\rangle \in \mathscr{H}_{\mathcal{R}}$ and $|\phi^{\mathcal{E}}_0\rangle \in \mathscr{H}_{\mathcal{E}}$ [37]. In fact, this type of initial state is often considered in the decoherence theory.

Summarizing the above discussions and taking into account the fact that the projection operators $\mathcal{P}_\mu$ are used in Part II of the third assumption when the condition C_D is satisfied, we propose the following definition for R-observable.

- A R-observable corresponds to an operator $A_{\{\mu\}}$ satisfying the following requirement: For all initial vectors with product form, $|\Psi(0)\rangle = |\psi^{\mathcal{R}}_0\rangle |\phi^{\mathcal{E}}_0\rangle$, there exists a decoherence time τ_d, such that if the condition C_D is satisfied for a time period $\mathcal{T} = [0, T]$ with $T > \tau_d$, then,

  $$\mathcal{P}_\mu \rho^{re}_{\mathcal{R}}(t) \mathcal{P}_\nu \doteq 0, \quad \forall \mu \neq \nu, \tag{8}$$

  for $t \in (\tau_d, T]$.

The decoherence time τ_d is usually a function of the value(s) of μ related to the initial condition $|\psi^{\mathcal{R}}_0\rangle$. Thus, more explicitly, one may write $\tau_d(\{\mu\})$.

A remark: Here, the decoherence time τ_d is defined by the requirement $\|\mathcal{P}_\mu \rho^{re}_{\mathcal{R}}(t) \mathcal{P}_\nu\| < \epsilon_x$. It is not exactly the same as the decoherence time τ^u_d usually discussed, which is defined by a decay to $1/e$ of the initial value. It is easy to verify the relation $\tau_d \sim -\tau^u_d \ln \epsilon_x$.

It is important to note that Eq.(8) is required to hold for *all* initial product vectors satisfying the condition C_D for a time period $\mathcal{T}$. This implies that the existence of a R-observable is independent of the concrete status of the environment under the specified condition. In this sense, the R-observable can be regarded as the system's 'own' property.

3. Fine and coarse-grained R-observables

We can show that a system $\mathcal{R}$ has a finest R-observable, such that all its other R-observables are coarse-grainings of the finest one. Let us first give the definition for a coarse-graining of a R-observable $A_{\{\mu\}}$. For this purpose, we arrange the labelling μ into groups labelled by η, such that each μ belongs to one and only one group η. Then, we can define coarse-grained projection operators $\mathcal{P}_\eta$ as

$$\mathcal{P}_\eta = \sum_{\mu \in \eta} \mathcal{P}_\mu, \tag{9}$$

and correspondingly an coarse-grained observable $A_{\{\eta\}}$,

$$A_{\{\eta\}} = \sum_\eta f(\eta) \mathcal{P}_\eta, \tag{10}$$

where $f(\eta)$ is a one-to-one mapping of η. It is easy to verify that, if Eq. (8) is satisfied by the projection operators $\mathcal{P}_\mu$, it is also satisfied by the coarse-grained projection operators $\mathcal{P}_\eta$. Hence, $A_{\{\eta\}}$ is also a R-observable.

Next, we show the existence of a finest R-observable. The key point is to note that here we consider a fixed set of reduced density matrices, which we denote by G_{re}, corresponding to a fixed set of initial conditions (product vectors) and to some times ($t \in (\tau_d, T]$). Obviously, there exists a finest division of the Hilbert space $\mathscr{H}_{\mathcal{R}}$ into orthogonal subspaces, such that all the matrices in G_{re} are block-diagonal with respect to this division. We use $\mathcal{P}_{\mu_f}$ to indicate the projection operators corresponding to this finest division; correspondingly we have an observable $A_{\{\mu_f\}}$ defined by Eq.(6). Clearly, by this construction, the observable $A_{\{\mu_f\}}$ satisfies the definition of R-observable. Furthermore, since the projection operators $\mathcal{P}_{\mu_f}$ give the finest division of $\mathscr{H}_{\mathcal{R}}$, all other R-observables must be coarse-grainings of $A_{\{\mu_f\}}$. Therefore, $A_{\{\mu_f\}}$ is the finest R-observable of the system $\mathcal{R}$. This completes our proof.

C. Explicit form of the condition C_D

In this section, we discuss the condition C_D. According to Part II of the general form of the third assumption, the condition C_D has the following meaning. That is, for an initial state of the total system described by $|\Psi(0)\rangle$, if this condition is satisfied for the time period $\mathcal{T} = [0, T]$, then, the state of the total system at the time $t = T$ has a mixed-state description $\rho(T)$ in Eq. (7).

Making use of arguments similar to those given in Sec. IV B 2 for the implications of definiteness, we see that Eq.(8) must hold for $t = T$. This requirement can not be fulfilled, if $\|\mathcal{P}_\nu H \mathcal{P}_\mu |\Psi(t)\rangle\|$ with $\nu \neq \mu$ is not negligibly small for some period of time before $t = T$. In fact, if there had been non-negligible transition among the subspaces $\mathscr{H}_\mu$ before $t = T$, usually the interaction might generate non-negligible elements of $\mathcal{P}_\mu \rho^{re}_{\mathcal{R}}(t) \mathcal{P}_\nu$ with $\mu \neq \nu$. Therefore, the condition C_D should include the following contents as a necessary part, that is, for certain time period before $t = T$, there is negligible transition among the subspaces $\mathscr{H}_\mu$.

Since Eq. (8) in fact represents a decoherence effect and it usually takes the decoherence time τ_d for a decoherence effect to happen, usually the above-discussed time period before $t = T$ should not be shorter than the decoherence time τ_d. Therefore, generally, the condition C_D requires negligible transition among the subspaces $\mathscr{H}_\mu$ for a time period $\mathcal{T} = [0, T]$ with $T \geq \tau_d$. Writing this explicitly, we have

$$\frac{1}{\langle\Psi|\Psi\rangle^{1/2}} \|\mathcal{P}_{\overline{\mu}} U(t,0) \mathcal{P}_\mu |\Psi(0)\rangle\| \doteq 0, \ \forall \mu, \ t \in \mathcal{T}, \quad (11)$$

where $\mathcal{P}_{\overline{\mu}} \equiv I - \mathcal{P}_\mu$. We call Eq. (11) the *non-transition condition* for $|\Psi(t)\rangle$ with respect to the R-observable $A_{\{\mu\}}$. Note that the non-transition condition also implies stableness of the corresponding property of $\mathcal{R}$. We do not see any other element that is necessary for the condition C_D to include, therefore, we assume that the non-transition condition can be taken as the condition C_D.

D. Assumption of mixed-state description

Now, we are ready to propose an explicit form of Part II of the third assumption. Before doing this, we need to determine the explicit form of the mixed-state description ρ in the general form of third assumption,

Let us first consider a description given by the internal observer, for which Part II of the third assumption is applicable. Suppose the non-transition condition is satisfied by $|\Psi(t)\rangle$ for a time period $\mathcal{T} = [0, T]$. According to the general form of the third assumption, this implies that at the time $t = T$ the total system is described by a vector $|\Psi_\mu(T)\rangle$ with a probability p_μ.

From the viewpoint of an imaginary external observer, according to the temporary assumption $\mathrm{A_T}$, the standard formalism of quantum mechanics is usable here, predicting that if a measurement is performed on the observable $A_{\{\mu\}} \otimes I_{\mathcal{E}}$, there is a probability $\langle\Psi(T)|\mathcal{P}_\mu|\Psi(T)\rangle / \langle\Psi(T)|\Psi(T)\rangle$ for a value μ to come out, with the system $\mathcal{R} + \mathcal{E}$ in a state described by $\mathcal{P}_\mu |\Psi(T)\rangle$. For the consistency of the two predictions given by the internal and the external observers, we get

$$|\Psi_\mu(T)\rangle = \frac{\mathcal{P}_\mu |\Psi(T)\rangle}{\|\mathcal{P}_\mu |\Psi(T)\rangle\|}, \ p_\mu = \frac{\langle\Psi(T)|\mathcal{P}_\mu|\Psi(T)\rangle}{\langle\Psi(T)|\Psi(T)\rangle}. \quad (12)$$

Thus, the explicit form of the mixed-state description $\rho(T)$ in Eq. (7) is obtained.

Based on the above discussions and summarizing results obtained in the previous sections, now we give an explicit form of Part II of the third assumption, which we term the assumption of mixed-state description.

- **Assumption of mixed-state description (MsD)**: If the total system $\mathcal{R} + \mathcal{E}$ has a description $|\Psi(t)\rangle$ that satisfies the non-transition condition (11) for a R-observable $A_{\{\mu\}}$ within a time period $\mathcal{T} = [0, T]$ with $T \geq \tau_d$, then, at the time T, with a probability $p_\mu = \frac{1}{\langle\Psi(T)|\Psi(T)\rangle} \langle\Psi(T)|\mathcal{P}_\mu|\Psi(T)\rangle$, the total system is described by $\mathcal{P}_\mu |\Psi(T)\rangle$.

We emphasize again that application of the assumption of MsD is subject to Part I of the third assumption.

Thus, at the time T, in addition to the Schrödinger evolution $|\Psi(T)\rangle$, the total system also has the following mixed-state description,

$$\rho(T) = \frac{1}{\langle\Psi(T)|\Psi(T)\rangle} \sum_\mu \mathcal{P}_\mu |\Psi(T)\rangle\langle\Psi(T)| \mathcal{P}_\mu. \quad (13)$$

Satisfaction of the non-transition condition implies that $\mathcal{P}_\mu |\Psi(t)\rangle \doteq U(t,0) \mathcal{P}_\mu |\Psi(0)\rangle$ for $t \in \mathcal{T}$.

A remark: Most of the contents in the assumption of MsD are given based on our experiences obtained in labs and an appropriate part of the standard formalism of quantum mechanics. The part lacking such a sound basis is the assumption that the non-transition condition is sufficient for the condition C_D. We'll give further discussions in this point, as well as the possibility of experimental test, in Sec. V E.

V. A CONCISE EXPRESSION OF THE THIRD ASSUMPTION

In this section, we derive an explicit expression of the principle of compatible description and obtain a mathematical expression for Part I of the third assumption. Making use of this result, a concise form of the third assumption can be obtained. We also give further discussion for the non-transition condition and for the possibility of experimental test.

A. Tree structure of branching for time evolution

Before deriving an explicit expression for the principle of compatible description, we first need to discuss a tree structure formed by the components of the mixed-state descriptions predicted by the assumption of MsD. For this purpose, we do not need to consider Part I of the third assumption. While, we must keep in mind that, only in the case that Part I of the third assumption is satisfied, could the branching picture given below be related to a valid time evolution.

Let us consider an initial normalized vector $|\Psi(t_0)\rangle$. According to the postulate of SE, for a time $t > t_0$ the state of the total system has a pure-vector description given by Schrödinger equation, $|\Psi(t)\rangle = U(t, t_0)|\Psi(t_0)\rangle$. Suppose the non-transition condition (11) is satisfied for a R-observable $A_{\{\mu_{(1)}\}}$ within a time interval $[\tau_1, \widetilde{\tau}_1]$, with $\widetilde{\tau}_1 - \tau_1 \geq \tau_d$. Then, according the assumption of MsD, for the time $t_1 = \tau_1 + \tau_d$, besides the pure-vector description $|\Psi(t_1)\rangle$, the total system also has the following mixed-state description [see Eq.(13)],

$$\rho(t_1) = \sum_{\mu_{(1)}} \mathcal{P}_{\mu_{(1)}} |\Psi(t_1)\rangle\langle\Psi(t_1)| \mathcal{P}^{\dagger}_{\mu_{(1)}}. \tag{14}$$

Each component in this mixed-state description evolves obeying Schrödinger equation, hence,

$$\rho(t) = \sum_{\mu_{(1)}} |\Psi_{(\mu_{(1)})}(t)\rangle\langle\Psi_{(\mu_{(1)})}(t)|, \quad \text{for } t > t_1, \tag{15}$$

where

$$|\Psi_{(\mu_{(1)})}(t)\rangle = U(t, t_1)\mathcal{P}_{\mu_{(1)}}|\Psi(t_1)\rangle. \tag{16}$$

For brevity, one may say that the vector $|\Psi(t)\rangle$ "*splits*" into the components $|\Psi_{(\mu_{(1)})}(t)\rangle$ at $t = t_1$. This feature is schematically plotted in Fig. 2.

Note that, although $|\Psi_{(\mu_{(1)})}(t)\rangle \in \mathscr{H}_{\mu_{(1)}}$ for $t \in (t_1, \widetilde{\tau}_1)$, beyond the time $\widetilde{\tau}_1$, it is not necessary for the vector $|\Psi_{(\mu_{(1)})}(t)\rangle$ to lie in the subspace $\mathscr{H}_{\mu_{(1)}}$, since the non-transition condition is not satisfied beyond $\widetilde{\tau}_1$. For this reason, in the subscript of Ψ we write $\mu_{(1)}$ in a pair of parentheses.

Suppose for a component $|\Psi_{(\mu_{(1)})}(t)\rangle$ of $t > \widetilde{\tau}_1$, the non-transition condition (11) is satisfied for a R-observable $A_{\{\mu_{(2)}\}}$ within a time period $[\tau_2, \widetilde{\tau}_2]$, with $\widetilde{\tau}_2 - \tau_2 \geq \tau_d$. Then, $|\Psi_{(\mu_{(1)})}(t)\rangle$ may split at the time $t_2 = \tau_2 + \tau_d$, giving the following mixed-state description according to the assumption of MsD,

$$\rho(t) = \sum_{\mu_{(2)}} |\Psi_{(\mu_{(1)}\mu_{(2)})}(t)\rangle\langle\Psi_{(\mu_{(1)}\mu_{(2)})}(t)|, \quad \text{for } t > t_2, \tag{17}$$

where

$$|\Psi_{(\mu_{(1)}\mu_{(2)})}(t)\rangle = U(t, t_2)\mathcal{P}_{\mu_{(2)}}|\Psi_{(\mu_{(1)})}(t_2)\rangle\rangle. \tag{18}$$

These features are also plotted in Fig. 2, where components like $|\Psi_{(\mu_{(1)})}(t)\rangle$ are indicated by short lines.

Proceeding with the above procedure, with increasing time, splitting of components may happen again and again. Since subscripts of Ψ will become even longer, for brevity, we use α to indicate a sequence of splittings and call it a *path* of splitting. Explicitly, we have

$$\alpha = \left(\mu^{\alpha}_{(1)}(t^{\alpha}_1) \rightarrow \mu^{\alpha}_{(2)}(t^{\alpha}_2) \rightarrow \ldots \rightarrow \mu^{\alpha}_{(n)}(t^{\alpha}_n)\right) \tag{19}$$

for a path of n splittings. For example, for $n = 2$ we have $|\Psi_{\alpha}(t) = |\Psi_{(\mu^{\alpha}_{(1)}\mu^{\alpha}_{(2)})}(t)\rangle$. Along a path α, around the i-th spitting, the non-transition condition (11) is satisfied for a R-observable $A_{\{\mu^{\alpha}_{(i)}\}}$ within a time period $[\tau^{\alpha}_i, \widetilde{\tau}^{\alpha}_i]$ with $\widetilde{\tau}^{\alpha}_i - \tau^{\alpha}_i \geq \tau_d$; the i-th spitting happens at the time $t^{\alpha}_i = \tau^{\alpha}_i + \tau_d$.



$|\Psi_{(\mu_{(1)}\mu_{(2)})}(t)>$
t_2
$|\Psi_{(\mu_{(1)}\mu'_{(2)})}(t)>$
$|\Psi_{(\mu_{(1)})}(t)>$
t_1
t_0
$|\Psi_{(\mu'_{(1)})}(t)>$
t_2
$|\Psi_{(\mu'_{(1)}\mu_{(2)})}(t)>$
$|\Psi_{(\mu'_{(1)}\mu'_{(2)})}(t)>$
t

FIG. 2: Schematic plot for the tree structure of time evolution. Each small square represents a splitting point of a component and each short line with direction represents Schrödinger evolution of a component (branch). Specifically, the tree plotted in this figure starts from an initial vector $|\Psi(t_0)\rangle$, then, evolves unitarily to a time t_1 and splits into components $|\Psi_{(\mu_{(1)})}(t_1)\rangle$. Then, after some period of unitary evolution, each component (branch) splits at a time t_2, giving a sub-branch $|\Psi_{(\mu_{(1)})(\mu_{(2)})}(t)\rangle$. A series of successive branches going from left to right form a path. Paths with different values of $\mu_{(1)}$ may have different times t_2 and different labeling $\mu_{(2)}$.

Several remarks about notations: (1) In the general situation here, a superscript α is added to μ, τ, and t belonging to a path α, since the values of τ_i, $\mu_{(i)}$, and t_i may be different along different paths α. (2) We write the number i in the subscript of $\mu^{\alpha}_{(i)}$ within parentheses, to indicate explicitly that the corresponding projection operators $\mathcal{P}_{\mu}$ at different splitting points i may be different. (3) A path α is in fact a function of the time t, hence, sometimes we write $\alpha(t)$.

The short lines and small squares in Fig. 2 form a structure like a tree. For this reason, we call one set of compatible splitting points and components, which stem from the same initial condition like those shown in Fig. 2, a *tree* and denote it by Υ. We call a component in a tree, represented by a short line in the figure, a *branch* of the tree.

Similar to Eqs. (16) and (18), it is easy to get the following explicit expression for a component at a time t, reached through a path α,

$$|\Psi_\alpha(t)\rangle = U(t,t_n^\alpha)\mathcal{P}_{\mu^\alpha_{(n)}}U(t_n^\alpha,t_{n-1}^\alpha)\mathcal{P}_{\mu^\alpha_{(n-1)}}\cdots U(t_{i+1}^\alpha,t_i^\alpha)\mathcal{P}_{\mu^\alpha_{(i)}}U(t_i^\alpha,t_{i-1}^\alpha)\cdots U(t_2^\alpha,t_1^\alpha)\mathcal{P}_{\mu^\alpha_{(1)}}U(t_1^\alpha,t_0)|\Psi(t_0)\rangle. \tag{20}$$

At the time t, the state of the total system can be described by both the vector $|\Psi(t)\rangle = U(t,t_0)|\Psi(t_0)\rangle$ and the following density operator,

$$\rho_\Upsilon(t) = \sum_{\alpha\in\Upsilon} |\Psi_\alpha(t)\rangle\langle\Psi_\alpha(t)|. \tag{21}$$

The probability for the realization of a path α is

$$P_\alpha(t) = \langle\Psi_\alpha(t)|\Psi_\alpha(t)\rangle, \tag{22}$$

with

$$\sum_\alpha \langle\Psi_\alpha(t)|\Psi_\alpha(t)\rangle = 1. \tag{23}$$

Using the relation $\sum_{\mu^\alpha_{(i)}} \mathcal{P}_{\mu^\alpha_{(i)}} = I$ for each pair (α, i), it is easy to verify that

$$|\Psi(t)\rangle = \sum_{\alpha\in\Upsilon} |\Psi_\alpha(t)\rangle. \tag{24}$$

B. Fine and coarse-grained trees

Starting from the same initial normalized vector $|\Psi(t_0)\rangle$, there may exist many trees. To see this point, it would be sufficient to note the following two points. That is, (i) the i-th splitting time t_i^α along a path α may in fact take any value between $\tau_i^\alpha + \tau_d$ and $\widetilde{\tau}_i^\alpha$; (2) if the non-transition condition is satisfied for one R-observable, then, it is also satisfied for a coarse-graining of the R-observable.

In this section, we show that the principle of compatible description and the existence of the finest R-observable (see Sec.IV B 3) require the existence of a finest tree among the trees starting from the same initial condition. Specifically, let us use Υ_f to denote a tree starting from an initial condition $|\Psi(t_0)\rangle$, which is obtained by taking a R-observable as fine as possible at each possible splitting time. Below, we prove that this tree Υ_f is the finest tree starting from $|\Psi(t_0)\rangle$.

Let us consider an arbitrarily chosen tree Υ, which has the first splitting point at a time t_1 for a R-observable $A_{\{\mu\}}$. Thus, at t_1, with the probability $p_\alpha(t_1) = \langle\Psi(t_1)|\mathcal{P}_\mu|\Psi(t_1)\rangle$, the total system $\mathcal{R}+\mathcal{E}$ is described by [see Eq.(14)]

$$|\Psi_\alpha(t_1)\rangle = \mathcal{P}_\mu|\Psi(t_1)\rangle \tag{25}$$

with a definite value of μ, where α is just μ in this case. According to the definition of Υ_f, the first splitting time of Υ_f is either t_1 or a time before t_1. Let us consider the latter case, since generalization of the following treatment to the former case is straightforward. Then, according to Υ_f, for times t just before t_1, the total system is described by a mixed-state description with components $|\Psi_{\alpha_f}(t)\rangle$.

To see what happens to the tree Υ_f at the time t_1, we note that, for Υ_f to give predictions compatible with those of Υ, as required by the principle of compatible description, each component $|\Psi_{\alpha_f}(t)\rangle$ must be able to split at the time t_1 into sub-components with definite values of μ. (The case that $|\Psi_{\alpha_f}(t)\rangle$ already has a definite value of μ can be treated in a similar way.) This implies that, within some appropriate time period before t_1, $|\Psi_{\alpha_f}(t)\rangle$ must satisfy the non-transition condition for the R-observable $A_{\{\mu\}}$, or for some R-observable(s) finer than $A_{\{\mu\}}$; we use $A_{\{\nu\}}$ to denote the finest one among these R-observables for the component $|\Psi_{\alpha_f}(t)\rangle$. Obviously, each subspace $\mathscr{H}_\nu$ is contained in some subspace $\mathscr{H}_\mu$, $\mathscr{H}_\nu \subseteq \mathscr{H}_\mu$.

Clearly, the (locally) finest R-observable $A_{\{\nu\}}$ generates the splitting of the component $|\Psi_{\alpha_f}(t)\rangle$ in the tree Υ_f at the time t_1. We use β_f to denote a newly generated path in the tree Υ_f at the time t_1. Thus, at the time t_1, in addition to the descriptions discussed above, the total system also has the following description, that is, with the probability $\langle\Psi_{\beta_f}(t_1)|\Psi_{\beta_f}(t_1)\rangle$, it is described by

$$|\Psi_{\beta_f}(t_1)\rangle = \mathcal{P}_\nu|\Psi_{\alpha_f}(t_1)\rangle. \tag{26}$$

The component $|\Psi_{\beta_f}(t_1)\rangle$ lies in the subspace $\mathscr{H}_\mu$ that contains the subspace $\mathscr{H}_\nu$, hence,

$$\mathcal{P}_\mu|\Psi_{\beta_f}(t_1)\rangle = |\Psi_{\beta_f}(t_1)\rangle. \tag{27}$$

Substituting Eq. (24) for Υ_f at the time t_1 into Eq. (25), then, making use of Eq. (27), we find

$$|\Psi_\alpha(t_1)\rangle = \sum_{\beta_f\in g_\alpha(t_1)} |\Psi_{\beta_f}(t_1)\rangle, \tag{28}$$

where $g_\alpha(t_1)$ is the set of paths β_f for which $|\Psi_{\beta_f}(t_1)\rangle$ lies in the same subspace $\mathscr{H}_\mu$ as $|\Psi_\alpha(t_1)\rangle$ does, that is,

$$g_\alpha(t_1) = \{\beta_f : |\Psi_{\beta_f}(t_1)\rangle \in \mathscr{H}_\mu, \ \ |\Psi_\alpha(t_1)\rangle \in \mathscr{H}_\mu\}. \tag{29}$$

In the above discussions, $A_{\{\mu\}}$ is the R-observable corresponding to the splitting point at the time t_1 along the path α. It is straightforward to generalize the above results to the case that $A_{\{\mu\}}$ is a coarse-graining of the R-observable at the time t_1 along the path α. In this general case, $A_{\{\mu\}}$ is not determined by $\alpha(t_1)$, hence, we write $g^\mu_\alpha(t_1)$ for the set in Eq.(29).

Beyond the time t_1, all the components $|\Psi_\alpha(t)\rangle$ and $|\Psi_{\beta_f}(t)\rangle$ evolves unitarily. For times t before the next splitting time of Υ, the tree Υ_f may have some splitting point(s). Since each splitting can be obtained by inserting the identity operator expressed as the summation of a complete set of projection operators [see Eq.(20)], an equation like Eq. (28) is still valid for these times t. Furthermore, the principle of compatible description requires that if $|\Psi_\alpha(t)\rangle \in \mathscr{H}_\mu$ for an arbitrary R-observable $A_{\{\mu\}}$, then, $|\Psi_{\beta_f}(t)\rangle \in \mathscr{H}_\mu$. A set g^μ_α can also be defined accordingly.

To study the influence of the next splitting point of Υ, we note that for times $t \geq t_1$ the components $|\Psi_\alpha(t)\rangle$ in the mixed-state description given by the tree Υ can be treated independently. Thus, we can treat each component $|\Psi_\alpha(t)\rangle$ of $t \geq t_1$ in the same way as that for the whole vector $|\Psi(t)\rangle$ of $t \geq 0$ discussed above. Similarly, we find that an equation similar to Eq. (28) holds for the time(s) t_2 with appropriately defined set(s) $g^\mu_\alpha(t)$.

Proceeding with the above procedure, finally, we get the following conclusion for an arbitrary time t: For each $\alpha(t) \in \Upsilon$, there exists a corresponding division of the set $\{\beta_f(t) \in \Upsilon_f\}$ into subsets $g^\mu_\alpha(t)$, such that

$$|\Psi_\alpha(t)\rangle = \sum_{\beta_f \in g^\mu_\alpha(t)} |\Psi_{\beta_f}(t)\rangle \quad \text{with} \\ |\Psi_{\beta_f}(t)\rangle \in \mathscr{H}_\mu \text{ if } |\Psi_\alpha(t)\rangle \in \mathscr{H}_\mu. \qquad (30)$$

Since $g^\mu_\alpha(t)$ changes only at splitting points, we can use the following method to determine the set $g^\mu_\alpha(t)$ for an arbitrary time. That is, (i) at a splitting point of Υ_f that does not coincide with a splitting point of Υ, since $|\Psi_\alpha(t)\rangle$ is not changed, the change of $g^\mu_\alpha(t)$ is directly given by the changes of the paths in Υ_f. (ii) At a splitting point of Υ, we can use the method of obtaining Eq. (29) to determine the change of the set $g^\mu_\alpha(t)$.

The relation given in Eq. (30) shows that Υ can be regarded as a coarse-graining of Υ_f. Therefore, due to the arbitrariness of Υ, Υ_f is the finest tree starting from the given initial condition. This completes our proof. In the appendix A, we show that it is not necessary for a coarse-graining at one splitting point of Υ_f to give a valid tree of paths.

C. A requirement of the principle of compatible description

In this section, we derive a requirement of the principle of compatible description. Suppose a tree Υ with paths α predicts that the system $\mathcal{R}$ may have some definite value μ of a R-observable $A_{\{\mu\}}$ at a time t. This implies that, for each component $|\Psi_\alpha(t)\rangle$, there exists one and only one value of μ which we denote by μ_α, such that

$$|\Psi_\alpha(t)\rangle \in \mathscr{H}_{\mu_\alpha}. \qquad (31)$$

For a given value of μ, we use s_μ to denote the set of paths α for which $\mu_\alpha = \mu$, i.e., $s_\mu = \{\alpha : |\Psi_\alpha(t)\rangle \in \mathscr{H}_\mu\}$. The component $|\Psi_\alpha(t)\rangle$ predicts that the system $\mathcal{R}$ has the probability [see Eq.(22)]

$$P_\Upsilon(\mu, \alpha, t) = \langle \Psi_\alpha(t)|\mathcal{P}_\mu|\Psi_\alpha(t)\rangle \qquad (32)$$

to possess a definite value μ. Thus, the probability $P_\Upsilon(\mu, t)$ for the system $\mathcal{R}$ to have the value μ at the time t, given by $\sum_\alpha P_\Upsilon(\mu, \alpha, t)$, has the following expression,

$$P_\Upsilon(\mu, t) = \sum_{\alpha \in s_\mu} \langle \Psi_\alpha(t)|\Psi_\alpha(t)\rangle. \qquad (33)$$

As discussed in the previous section, there exists a finest mixed-state description given by the finest tree Υ_f. According to Eq. (30), as a result of the relation (31), each component $|\Psi_{\beta_f}(t)\rangle$ lies in a subspace with a definite value of μ, specifically, $|\Psi_{\beta_f}(t)\rangle \in \mathscr{H}_{\mu_\alpha}$ for $\beta_f \in g_\alpha(t)$. Let us use r_μ to denote the set of paths β_f for which the corresponding component lies in the subspace $\mathscr{H}_\mu$, that is, $r_\mu := \{\beta_f : |\Psi_{\beta_f}(t)\rangle \in \mathscr{H}_\mu\}$. Then, the finest tree Υ_f predicts the following probability for the system $\mathcal{R}$ to have a definite value μ,

$$P_{\Upsilon_f}(\mu, t) = \sum_{\beta_f \in r_\mu} \langle \Psi_{\beta_f}(t)|\Psi_{\beta_f}(t)\rangle. \qquad (34)$$

Making use of the relations given in Eq.(30), it is not difficult to verify the following relation between the sets r_μ, s_μ, and $g^\mu_\alpha(t)$,

$$r_\mu = \bigcup_{\alpha \in s_\mu} g^\mu_\alpha(t). \qquad (35)$$

Substituting Eq. (30) into Eq. (33), we get

$$P_\Upsilon(\mu, t) = \sum_{\alpha \in s_\mu} \sum_{\beta_f, \beta'_f \in g^\mu_\alpha(t)} \langle \Psi_{\beta_f}(t)|\Psi_{\beta'_f}(t)\rangle. \qquad (36)$$

The principle of compatible description requires that $P_\Upsilon(\mu, t) = P_{\Upsilon_f}(\mu, t)$. Comparing Eq.(34) with Eq.(36) and making use of the relation given in Eq. (35), we obtain

$$\sum_{\alpha \in s_\mu} \sum_{\beta_f \neq \beta'_f \in g^\mu_\alpha} \langle \Psi_{\beta_f}(t)|\Psi_{\beta'_f}(t)\rangle \doteq 0, \quad \forall \mu. \qquad (37)$$

Without a complete proof, we conjecture that the validity of Eq. (37) for all the trees Υ and for all the times t implies the following relation,

$$\mathcal{D}_{\beta_f \beta'_f} \doteq \delta_{\beta_f \beta'_f} \mathcal{D}_{\beta_f \beta_f}, \qquad \forall \beta_f, \beta'_f \in \Upsilon_f, \qquad (38)$$

where

$$\mathcal{D}_{\beta_f \beta'_f} := \langle \Psi_{\beta_f}(t)|\Psi_{\beta'_f}(t)\rangle. \qquad (39)$$

Taking this conjecture, the principle of compatible description requires Eqs.(30) and (38) to hold.

Although we do not have a complete proof for the above conjecture, we do have arguments for its correctness. In fact, for Eq. (37) to hold for an arbitrary tree Υ, generally it is reasonable to expect that $\mathrm{Re}\langle \Psi_{\beta_f}(t)|\Psi_{\beta'_f}(t)\rangle \doteq 0$ for $\beta_f \neq \beta'_f$. Then, since $\langle \Psi_{\beta_f}(t)|\Psi_{\beta'_f}(t)\rangle$ is a dynamical quantity that changes with Schrödinger evolution except at splitting points, if its real part vanishes for all the times, usually its imaginary part should vanish as well. The difficulty in completing the proof for Eq. (38) is related to the fact that mere coarse-graining of an arbitrary splitting point of Υ_f does not necessarily give a valid tree (see the appendix A), although each Υ is a coarse-graining of Υ_f for the same initial condition [see Eq.(30)]. The missing part of the proof is to show that this restriction to the construction of valid trees by the procedure of coarse-graining does not influence Eq. (38).

D. Initial-vector-restriction and a concise expression of the third assumption

In this section, based on results obtained above, we give a concise expression of the third assumption. For this purpose, we first need to find an explicit mathematical expression for Part I of the third assumption, namely, a condition under which all the mixed-state descriptions given by trees starting from the same initial condition $|\Psi(t_0)\rangle$ give consistent predictions for the probabilities for $\mathcal{R}$ to have definite properties.

Below, we show that the above-mentioned condition takes the following form: Namely, for trees starting from $|\Psi(t_0)\rangle$, there exists a finest tree Υ_f such that both Eqs. (30) and (38) hold. Under this condition, it is not difficult to show that the principle of compatible description is obeyed. Indeed, when Eq. (30) holds, the probability $P_\Upsilon(\mu,t)$ is given by Eq. (36). Then, substituting Eq. (38) into Eq. (36), we have

$$P_\Upsilon(\mu,t) = \sum_{\alpha\in s_\mu}\sum_{\beta_f\in g^\mu_\alpha(t)} P_{\beta_f}(t) = \sum_{\beta_f\in r_\mu} P_{\beta_f}(t), \qquad (40)$$

where $P_{\beta_f}(t) = \langle\Psi_{\beta_f}(t)|\Psi_{\beta_f}(t)\rangle$ is the probability for the realization of the path β_f in the tree Υ_f. Therefore, the probability $P_\Upsilon(\mu,t)$ is given by the corresponding probabilities $P_{\beta_f}(t)$ according to the sum rule of probability. Making use of this result, it is not difficult to see that all the mixed-state descriptions give consistent predictions for the probabilities for the system $\mathcal{R}$ to have definite values of μ.

It is straightforward to generalize the above discussions to the case that a fraction of the paths of a tree Υ give components possessing definite values of μ. The result is similar: The principle of compatible description is satisfied, if the two equations (30) and (38) hold.

Finally, we get the following concise expression of the third assumption, without explicitly mentioning the principle of compatible description.

- **The third assumption**: The assumption of MsD is applicable to vectors selected by the Initial-vector-restriction.

Here, the Initial-vector-restriction is defined as follows.

- **Initial-vector-restriction**: Only those initial vectors are considered, for each of which there exists a finest tree Υ_f that satisfies Eq.(38) and has the relation in Eq.(30) to all other trees starting from the same initial condition.

Below we give several remarks concerning the Initial-vector-restriction.

(1) If a vector $|\Psi_0\rangle$ does not satisfy the Initial-vector-restriction, then, at least, it can not describe a state of the total system $\mathcal{R}+\mathcal{E}$, in which $\mathcal{R}$ can be used as the essential part of a measuring apparatus.

It is of interest to consider the extreme case that a vector $|\Psi_0\rangle$ satisfies the Initial-vector-restriction for none of the subsystems of the total system. If the total system could lie in a state described by such a vector $|\Psi_0\rangle$, there would exist no internal measuring apparatus, hence, no possibility of experimentally checking predictions of this vector $|\Psi_0\rangle$ from inside. This would be physically meaningless for a total system without any external system (like the universe). In this case, it seems reasonable to assume that such a vector $|\Psi_0\rangle$ does not describe any physical state of the total system.

(2) Physical restriction to the initial condition is not a new idea in physics, in particular, when dealing with irreversible processes. In fact, in studying the microscopic origin of the macroscopic irreversibility stated in the second law of thermodynamics, it has been suggested by many authors that there might exist some selection rule for the initial condition (see, e.g., [38, 39]). Indeed, only for certain types of initial conditions, have master equations been derived for open quantum systems [39, 40].

(3) In case that Eq. (38) can not be rigorously derived from Eq. (37), we suggest that the Initial-vector-restriction should be used as Part I of the third assumption, because it can guarantee satisfaction of the principle of compatible description and it has an explicit, mathematical expression.

(4) This Initial-vector-restriction is irrelevant to the restriction used in the definition of R-observable, which states that only initial vectors of the product form are considered. In fact, as discussed in Sec.IV B 2, the restriction considered in the definition of R-observable has nothing to do with the present physical state of the total system under consideration. The purpose of considering initial product vectors in the definition of R-observable is just to determine the subspaces $\mathscr{H}_{\mathcal{R}\mu}$, such that decoherence may happen with respect to them, to capture the observed initial-state-independent feature of definite properties of measuring apparatuses. By contrast, the Initial-vector-restriction is required by the principle of compatible description and it selects physical-state vectors for which the assumption of MsD is applicable.

In including this section, we show that the temporary assumption $\mathrm{A_T}$ given in Sec.IV A can be derived from the theory proposed above. Let us consider the big system composed of the external observer and $\mathcal{R}+\mathcal{E}$. Since there is no interaction between the external observer and $\mathcal{R}+\mathcal{E}$, we assume that the big system is initially described by $|\Psi(t_0)\rangle \otimes |\xi_0\rangle$, where $|\xi_0\rangle$ is a vector in the Hilbert space of the external observer. Suppose the vector $|\Psi(t_0)\rangle$ satisfies the Initial-vector-restriction for the system $\mathcal{R}+\mathcal{E}$ and application of the assumption of MsD gives paths $|\Psi_\alpha(t)\rangle$. For the big system, application of the assumption of MsD will give paths $|\Psi_\alpha(t)\rangle \otimes |\xi(t)\rangle$. Here, since there is no interaction between the external observer and $\mathcal{R}+\mathcal{E}$, there is no correlation between the time evolution of the external observer and of $\mathcal{R}+\mathcal{E}$, hence, for the simplicity in discussion we use the same $|\xi(t)\rangle$ for all the paths α. Then, it is easy to verify that these paths for the big system also satisfy the Initial-vector-restriction, that is, they satisfy both Eqs.(30) and (38), therefore, the assumption of MsD is applicable to the vector $|\Psi(t)\rangle \otimes |\xi(t)\rangle$ for the big system.

It is not difficult to see that, if $|\Psi(t)\rangle$ satisfies the non-transition condition for a period $\mathcal{T}$, the same is true for $|\Psi(t)\rangle \otimes |\xi(t)\rangle$. Then, application of the assumption of MsD to the big system also predicts that the system $\mathcal{R}+\mathcal{E}$ has the mixed-state description given in Eq.(13). This is just what is stated in the temporary assumption $\mathrm{A_T}$.

E. Further discussions in the condition C_D and the possibility of experimental test

In this section, we give further discussions for taking the non-transition condition as the condition C_D in the general form of the third assumption (see Sec.IV C and Sec.IV D for previous discussions), as well as the possibility of experimental test for this choice of the condition C_D. It is important to note that Part I of the third assumption guarantees the physical consistency of the proposed theory, independent of the choice of the condition C_D.

In the theory proposed above, the non-transition condition alone can not guarantee the appearance of some definite property of the considered system $\mathcal{R}$. In fact, for the system $\mathcal{R}$ to possess a definite property, other (more important) requirements must also be met, namely, the principle of compatible description and the existence of R-observable. In particular, Eq.(38) imposes a restriction more stringent than Eq. (8), since the former must be satisfied in all the future times.

In fact, it is experiments that may finally determine whether the non-transition condition is sufficient for the condition C_D. To see this point, let us first discuss what may happen, if the non-transition condition is looser than what is really needed for the condition C_D, but it is still employed as the condition C_D in the third assumption. In this case, the assumption of MsD will predict more definite properties than the system $\mathcal{R}$ may really possess. In principle, such predictions can be tested experimentally, by studying possible coherence among the components of the mixed-state descriptions predicted by the assumption of MsD. Therefore, at least in principle, experiments may test whether the non-transition condition is looser than what is really needed for the condition C_D.

Next, let us discuss what may happen, if the non-transition condition is sufficient for the condition C_D, but in addition to it some further requirement is added to the condition C_D. In this case, the Initial-vector-restriction will have more requirements, as a result, there will be less vectors that may satisfy the Initial-vector-restriction. The more stringent the condition C_D is, the less the valid vectors will be (in the sense of satisfying the Initial-vector-restriction), as a result, the predictability and explainablility of the theory will be reduced. In this case, there may exist experimental results that can not be explained by the theory. For example, under a too stringent condition C_D, there might exist no vector that can satisfy the Initial-vector-restriction.

Therefore, pure theoretical considerations can not completely determine the exact form of the condition C_D, i.e., whether the assumed non-transition condition in the assumption of MsD is indeed sufficient or not. It is experiments that may finally determine the exact form of the condition C_D, hence, in principle, the proposed theory is experimentally testable.

VI. MEASURING APPARATUS AND MEASUREMENT

In this section, we use the theory proposed above to give a brief discussion for measuring apparatus and measurement process, in particular, for the contents of the axiom of measurement in the standard formalism of quantum mechanics.

Let us first give a definition for measuring apparatus in a general sense. If the total system lies in an initial state described by $|\Psi_0\rangle$ that satisfies the Initial-vector-restriction, then, according to the assumption of MsD, it is possible for the system $\mathcal{R}$ to have some definite properties within some time periods in the future. In principle, such a system $\mathcal{R}$ may be used as the essential part of a measuring apparatus; in the case that $\mathcal{R}$ is a microscopic system, some auxiliary system is needed to amplify information contained in $\mathcal{R}$.

Usually, a measurement should have the following two features: (i) The interaction between the measuring apparatus $\mathcal{R}$ and the measured system may induce transition among the subspaces $\mathscr{H}_\mu$ related to a R-observable $A_{\{\mu\}}$ of $\mathcal{R}$; and (ii) after a measurement process, the measuring apparatus has some definite value μ of the R-observable.

Next, we show that, for appropriately designed measurement schemes, it is possible to derive the contents of the axiom of measurement in the standard formalism of quantum mechanics. Let us consider a measurement on an observable B of a system $\mathcal{S}$, with normalized eigenvectors $|b\rangle$, $B|b\rangle = b|b\rangle$. Suppose the initial vector has a product form, satisfying the Initial-vector-restriction,

$$|\Psi(t_0)\rangle = \left(\sum_b C_b(t_0)|b\rangle \right) |R_0\rangle, \tag{41}$$

where $|R_0\rangle \in \mathscr{H}_\mathcal{R}$ is normalized. The measured system $\mathcal{S}$ is a part of the environment $\mathcal{E}$ of $\mathcal{R}$. For brevity, in this section, we do not explicitly mention other systems in the environment $\mathcal{E}$, though they do exist.

Schrödinger evolution of $|\Psi(t_0)\rangle$ in Eq. (41) can be written as

$$|\Psi(t)\rangle = \sum_b C_b(t)|b\rangle|R_b(t)\rangle, \tag{42}$$

with normalized $|R_b(t)\rangle$. Suppose the interaction process can be designed, such that, for a time τ_1, (i) each $|R_b(\tau_1)\rangle$ lies in a subspace $\mathscr{H}_{\mathcal{R}\mu}$ and can be written as $|R_{\mu(b)}(\tau_1)\rangle$, where μ is a function of b, with $\mu(b) \neq \mu(b')$ for $b \neq b'$, (ii) within a time interval $[\tau_1, t_1]$ with $t_1 - \tau_1 > \tau_d$, the non-transition condition (11) is satisfied. Then, at the time t_1, according to the assumption of MsD, in addition to the description $|\Psi(t)\rangle$, the composite system $\mathcal{R} + \mathcal{S}$ also has the following description: Namely, with a probability $|C_a(t_1)|^2 = |C_a(\tau_1)|^2$, it is described by $|b\rangle|R_{\mu(b)}(t_1)\rangle$, with $\mathcal{S}$ described by $|b\rangle$ and $\mathcal{R}$ possessing a definite value of μ. The value of b can be inferred from the value of μ obtained after the measurement, which in principle can be recorded (perhaps after some amplification process). Thus, we get the basic contents of the axiom of measurement in the standard formalism of quantum mechanics.

The above discussions show that there is no "collapse of state vector" in the theory proposed in this paper. Here, a mixed-state description ρ for the total system may appear, just because it may describe the same physical state of the total system as a pure-vector description $|\Psi(t)\rangle$ does.

VII. DECOHERENCE MECHANISM

In this section, we first discuss the time evolution of the total system when the non-transition condition is satisfied (Sec. VII A). Then, making use of the results obtained, we discuss the decoherence mechanism for the existence of R-observable (Sec. VII B) and for Eq. (38) to hold (Sec. VII C).

A. Time evolution under the non-transition condition

Let us first discuss the time evolution within a time period $\mathcal{T} = [0, T]$, in which the non-transition condition Eq.(11) is satisfied. This condition implies that $\mathcal{P}_{\overline{\mu}} H \mathcal{P}_\mu |\Psi(t)\rangle \doteq 0$, hence,

$$\mathcal{P}_\mu H |\Psi(t)\rangle \doteq \mathcal{P}_\mu H \mathcal{P}_\mu |\Psi(t)\rangle \quad \forall\ \mu,\ t \in \mathcal{T}. \tag{43}$$

That is, $\mathcal{P}_\mu$ is effectively commutable with H for the vectors $|\Psi(t)\rangle$ within this time period. Then, multiplying Schrödinger equation (1) by $\mathcal{P}_\mu$ from the left, we get the following equation of motion,

$$\mathrm{i}\hbar \frac{\partial}{\partial t} |\Psi_\mu(t)\rangle \doteq H_\mu |\Psi_\mu(t)\rangle \quad \text{for } t \in \mathcal{T}, \tag{44}$$

where $|\Psi_\mu(t)\rangle = \mathcal{P}_\mu |\Psi(t)\rangle$ and

$$H_\mu \equiv \mathcal{P}_\mu H \mathcal{P}_\mu = \mathcal{P}_\mu H_\mathcal{R} \mathcal{P}_\mu + \mathcal{P}_\mu H_I \mathcal{P}_\mu + H_\mathcal{E} \mathcal{P}_\mu. \tag{45}$$

Note that H_μ is an operator acting in the subspace $\mathscr{H}_{\mathcal{R}\mu} \otimes \mathscr{H}_\mathcal{E}$. Equation (44) has the following formal solution,

$$|\Psi_\mu(t)\rangle \doteq \exp\left(-\mathrm{i} H_\mu t/\hbar\right) |\Psi_\mu(0)\rangle \quad \text{for } t \in \mathcal{T}. \tag{46}$$

Next, we discuss an implication of the non-transition condition. The non-transition condition (11) for the time period $\mathcal{T}$ is in fact equivalent to the relation $\mathcal{P}_{\overline{\mu}} H \mathcal{P}_\mu |\Psi(t)\rangle \doteq 0$, hence, is equivalent to

$$\mathcal{P}_{\overline{\mu}} H_\mathcal{R} \mathcal{P}_\mu |\Psi(t)\rangle \doteq -\mathcal{P}_{\overline{\mu}} H_I \mathcal{P}_\mu |\Psi(t)\rangle \quad \forall \mu,\ t \in \mathcal{T}. \tag{47}$$

Note that the operator $\mathcal{P}_{\overline{\mu}} H_\mathcal{R} \mathcal{P}_\mu$ on the left hand side of Eq.(47) has trivial action in $\mathscr{H}_\mathcal{E}$, the Hilbert space of the environment $\mathcal{E}$, while the operator $\mathcal{P}_{\overline{\mu}} H_I \mathcal{P}_\mu$ on the right hand side has non-trivial action in $\mathscr{H}_\mathcal{E}$. Hence, generally, the non-transition condition requires that for all values of μ and for $t \in \mathcal{T}$,

$$\mathcal{P}_{\overline{\mu}} H_\mathcal{R} \mathcal{P}_\mu |\Psi(t)\rangle \doteq 0, \tag{48}$$

$$\mathcal{P}_{\overline{\mu}} H_I \mathcal{P}_\mu |\Psi(t)\rangle \doteq 0. \tag{49}$$

Note that an important case in which Eq.(48) is satisfied is that $\mathcal{P}_{\overline{\mu}} H_\mathcal{R} \mathcal{P}_\mu \doteq 0$, or equivalently,

$$[H_\mathcal{R}, A_{\{\mu\}}] \doteq 0. \tag{50}$$

In this case, the subspaces $\mathscr{H}_{\mathcal{R}\mu}$ are approximately eigen-subspaces of $H_\mathcal{R}$ and the corresponding definite property of $\mathcal{R}$ is stable as long as the interaction is weak.

B. Decoherence mechanism for the existence of R-observable

In this section, we show that Eq.(8) in the definition of R-observable, which is in fact a decoherence effect, can be written as a requirement in the property of a generalized (quantum) Loschmidt echo. Then, we discuss a condition under which an observable $A_{\{\mu\}}$ can be a R-observable.

To reveal the mechanism for Eq. (8) to hold, we express its left hand side in terms of quantities in $\mathscr{H}_\mathcal{E}$, similar to a method used in the study of preferred pointer states [48, 49]. For this purpose, let us consider an arbitrary basis in the subspace $\mathscr{H}_{\mathcal{R}\mu}$, which we denote by $|m_\mu\rangle \in \mathscr{H}_{\mathcal{R}\mu}$. In this basis, Eq. (8) has the following equivalent form,

$$\langle m_\mu | \mathrm{Tr}_\mathcal{E} \left[|\Psi(t)\rangle\langle\Psi(t)| \right] |n_\nu\rangle \doteq 0 \tag{51}$$

for $t \in [\tau_d, T]$ and for all m_μ and n_ν of $\mu \neq \nu$. Making use of the expression $|\Psi_\mu(t)\rangle = \mathcal{P}_\mu |\Psi(t)\rangle$ and the formal solution in Eq.(46), it is not difficult to find that Eq.(51) can be written as

$$\langle\Psi(0)| e^{\mathrm{i} H_\nu t/\hbar} |n_\nu\rangle\langle m_\mu| e^{-\mathrm{i} H_\mu t/\hbar} |\Psi(0)\rangle \doteq 0. \tag{52}$$

Let us consider initial vectors of the product form, as required in the definition of R-observable,

$$|\Psi(0)\rangle = \left(\sum_{\mu} \sum_{m_\mu} c_{m_\mu}(0) |m_\mu\rangle \right) |\phi_0\rangle, \tag{53}$$

with $|\phi_0\rangle \in \mathscr{H}_{\mathcal{E}}$. If the non-transition condition does not impose too stringent restriction to the coefficients $c_{m_\mu}(0)$, then, due to the arbitrariness of the coefficients $c_{m_\mu}(0)$, Eq.(52) is equivalent to the following requirement,

$$\langle \phi_0 | \langle n'_\nu | e^{\mathrm{i}H_\nu t/\hbar} | n_\nu \rangle \langle m_\mu | e^{-\mathrm{i}H_\mu t/\hbar} | m'_\mu \rangle | \phi_0 \rangle \doteq 0. \tag{54}$$

Finally, introducing the operator

$$V_{m_\mu m'_\mu}(t) := \langle m_\mu | e^{-\mathrm{i}H_\mu t/\hbar} | m'_\mu \rangle, \tag{55}$$

which represents a non-unitary evolution in the Hilbert space of the environment $\mathcal{E}$, we find that Eq.(8) has the following form, namely,

$$L_G(t) \doteq 0 \tag{56}$$

for $t \in [\tau_d, T]$ and for all m_μ, m'_μ, n_ν and n'_ν of $\mu \neq \nu$, where

$$L_G(t) := \langle \phi_0 | V^\dagger_{n_\nu n'_\nu}(t) V_{m_\mu m'_\mu}(t) | \phi_0 \rangle \tag{57}$$

is the overlap two non-unitary evolutions in the Hilbert space of the environment.

The quantity $L_G(t)$ defined in Eq. (57) has a form similar to the so-called quantum Loschmidt-echo amplitude. To see this point, we recall that, as a measure of the stability of the quantum motion of a system under small perturbation, the quantum Loschmidt echo is defined as the overlap of the time evolution of the same initial state $|\psi(0)\rangle$ under two Hamiltonians H_0 and H_1 [41],

$$M(t) \equiv |m(t)|^2 = |\langle \psi(0) | e^{\mathrm{i}H_1 t/\hbar} e^{-\mathrm{i}H_0 t/\hbar} | \psi(0) \rangle|^2. \tag{58}$$

Decaying behaviors of the Loschmidt echo with small difference between H_0 and H_1 have been extensively studied in recently years (see review given in Ref.[42]). In particular, when the two systems H_0 and H_1 are quantum chaotic systems, or quantum integrable systems possessing classical counterparts with sufficiently large degrees of freedom, the echo has typically an exponential decay [42–47].

Loosely speaking, the following four factors are responsible for the decaying behavior of the Loschmidt echo $M(t)$. That is, (i) Schrödinger evolution of the two systems H_0 and H_1 start from the same initial condition; (ii) there exists some difference between H_0 and H_1; (iii) the Hilbert space is sufficiently large, such that there is no finite-dimension effect in the separation of the two trajectories (evolutions) in the Hilbert space. Indeed, when these three requirements are met, the two systems have different trajectories in the Hilbert space, separating with increasing time. If, furthermore, the following fourth requirement is met, i.e., (iv) at least one of the two systems has sufficiently irregular motion in the Hilbert space within the time period of interest, then, the Loschmidt echo may have a fast decay, usually, an exponential decay.

Similar arguments are also applicable to the quantity $L_G(t)$. Indeed, $L_G(t)$ is also an overlap of two evolutions starting from the same initial condition. The definition in Eq.(55) suggests that if the unitary evolution under H_μ is sufficiently irregular, the operators $V_{m_\mu m'_\mu}(t)$ may generate somewhat irregular motions in the Hilbert space of the environment. Then, if there are sufficient differences among the effects of H_μ and if the Hilbert space of $\mathcal{E}$ is sufficiently large, it is reasonable to expect that the quantity $|L_G(t)|$ has a fast decay, like the Loschmidt echo. In this case, Eq. (56) may hold, as a result, $A_{\{\mu\}}$ may be a R-observable.

To summarize, $A_{\{\mu\}}$ may be a R-observable, when the following requirements are met:

1. The operators $V_{m_\mu m'_\mu}(t)$ generate sufficiently irregular motion in the Hilbert space of the environment.
2. The effective Hamiltonians H_μ have sufficiently different influences in the motions generated by $V_{m_\mu m'_\mu}(t)$.
3. The Hilbert space of the environment $\mathcal{E}$ is sufficiently large.

C. $\mathcal{D}_{\alpha\alpha'}$ expressed as a generalized Loschmidt echo

In this section, we show that Eq. (38) may hold due to a mechanism similar to that discussed in the previous section. When Eq. (38) holds, as a result of Eq. (30), all other trees have a similar property, i.e.,

$$\mathcal{D}_{\alpha\alpha'} \doteq \delta_{\alpha\alpha'} \mathcal{D}_{\alpha\alpha}, \quad \forall \alpha, \alpha' \in \Upsilon \text{ and } \forall \Upsilon, \tag{59}$$

where

$$\mathcal{D}_{\alpha\alpha'} := \langle \Psi_\alpha(t) | \Psi_{\alpha'}(t) \rangle. \tag{60}$$

Below, we show that $\mathcal{D}_{\alpha\alpha'}$ can also be expressed in the form of a generalized Loschmidt echo.

For the above mentioned purpose, it would be more convenient to rewrite the component $|\Psi_\alpha(t)\rangle$ in Eq. (20) in a form with respect to the times τ_i and $\widetilde{\tau}_i$. The reason is that the non-transition condition (11) is satisfied within the time intervals $[\tau_i^\alpha, \widetilde{\tau}_i^\alpha]$. Making use of Eq. (46) for the time intervals $[\tau_i^\alpha, \widetilde{\tau}_i^\alpha]$, we can write $|\Psi_\alpha(t)\rangle$ in Eq. (20) in the following form,

$$|\Psi_\alpha(t)\rangle = W_\alpha(t, t_0) |\Psi(t_0)\rangle, \tag{61}$$

where

$$\begin{aligned} W_\alpha(t, t_0) = {} & U(t, \widetilde{\tau}_n^\alpha)\; U_{\mu^\alpha_{(n)}}(\widetilde{\tau}_n^\alpha, \tau_n^\alpha)\; U(\tau_n^\alpha, \widetilde{\tau}_{n-1}^\alpha) \\ & \cdots U_{\mu^\alpha_{(2)}}(\widetilde{\tau}_2^\alpha, \tau_2^\alpha) U(\tau_2^\alpha, \widetilde{\tau}_1^\alpha) U_{\mu^\alpha_{(1)}}(\widetilde{\tau}_1^\alpha, \tau_1^\alpha) U(\tau_1^\alpha, t_0), \end{aligned} \tag{62}$$

with the following definition of $U_{\mu^\alpha_{(i)}}(\widetilde{\tau}^\alpha_i, \tau^\alpha_i)$,

$$U_\mu(t',t) := \exp\left\{-\frac{\mathrm{i}}{\hbar}(t'-t)H_\mu\right\}. \tag{63}$$

In the derivation of Eq.(61), we have used the following property related to times $t^\alpha_i \in (\tau^\alpha_i, \widetilde{\tau}^\alpha_i)$, namely,

$$U(\widetilde{\tau}^\alpha_i, t^\alpha_i)\mathcal{P}_{\mu^\alpha_{(i)}} U(t^\alpha_i, \tau^\alpha_i) = U_{\mu^\alpha_{(i)}}(\widetilde{\tau}^\alpha_i, \tau^\alpha_i), \tag{64}$$

which can be obtained by making use of Eq. (43). For $t \in (\tau^\alpha_n, \widetilde{\tau}^\alpha_n)$, the operator $W_\alpha(t,t_0)$ can be obtained by replacing the first two terms on the right hand side of Eq. (62) by $U_{\mu^\alpha_{(n)}}(t, \tau^\alpha_n)$.

The operator $W_\alpha(t,t_0)$ generates a time evolution in the total Hilbert space, given by a sequence of unitary operators U in the total Hilbert space $\mathscr{H}$, separated by unitary operators U_μ acting in subspaces $\mathscr{H}_\mu$. As a product of Us and U_μs, $W_\alpha(t,t_0)$ is no longer a unitary operator in the total Hilbert space $\mathscr{H}$. Making use of Eq.(61), $D_{\alpha\alpha'}$ can be written as

$$D_{\alpha\alpha'} = \langle\Psi(t_0)|W^\dagger_\alpha(t,t_0)W_{\alpha'}(t,t_0)|\Psi(t_0)\rangle. \tag{65}$$

Obviously, like $L_G(t)$ discussed in the previous section, $D_{\alpha\alpha'}$ expressed in Eq.(65) can also be regarded as a generalized Loschmidt-echo amplitude, with unitary operators in Eq.(58) replaced by the operators W.

Then, arguments similar to those given in the previous section for $L_G(t)$ can be applied to the quantity $D_{\alpha\alpha'}$, as well, giving the following results: Equation (59) may hold, when the following requirements are met:

1. The operators $W_\alpha(t,t_0)$ generate sufficiently irregular motion in the total Hilbert space.
2. The differences among H_μ should be sufficiently large, such that $W_\alpha(t,t_0)$ of each two different paths generate sufficiently different motions in the Hilbert space.
3. The total Hilbert space is sufficiently large.

VIII. APPLICATION I — ISOLATABLE AND NON-ISOLATABLE SYSTEMS

In this section, as an application of the theory proposed above, we show that a system which is isolatable from its environment for a sufficiently long time period does not have a practically meaningful R-observable. We still use $\mathcal{R}$ to denote the considered system in this section, even though a system without a R-observable can not be used as a measuring apparatus.

Suppose there exists an initial normalized vector with a product form, $|\Psi(0)\rangle = |\psi^\mathcal{R}_0\rangle|\phi^\mathcal{E}_0\rangle$, for which the system $\mathcal{R}$ is isolated from its environment $\mathcal{E}$ for a practically-infinitely-long time period $\mathcal{T}$ in the following sense,

$$H_I|\Psi(t)\rangle \doteq 0 \quad \text{for } t \in \mathcal{T} = [0,T]. \tag{66}$$

Making use of Eq.(66), we find that for $t \in \mathcal{T}$,

$$e^{-\mathrm{i}Ht/\hbar}|\Psi(0)\rangle \doteq e^{-\mathrm{i}H_\mathcal{R}t/\hbar}|\psi^\mathcal{R}_0\rangle e^{-\mathrm{i}H_\mathcal{E}t/\hbar}|\phi^\mathcal{E}_0\rangle. \tag{67}$$

Making use of the definition of the reduced density matrix of the system $\mathcal{R}$, it is straightforward to verify that its elements in a basis $\{|m_\mu\rangle\}$ of the subspace $\mathscr{H}_{\mathcal{R}\mu}$ (used in Sec.VII B) can be expressed as

$$\langle m_\mu|\rho^{re}_\mathcal{R}(t)|n_\nu\rangle = \langle\Psi(0)|e^{\mathrm{i}Ht/\hbar}|n_\nu\rangle\langle m_\mu|e^{-\mathrm{i}Ht/\hbar}|\Psi(0)\rangle. \tag{68}$$

Substituting Eq.(67) into Eq.(68), one has

$$\langle m_\mu|\rho^{re}_\mathcal{R}(t)|n_\nu\rangle \doteq \langle\psi^\mathcal{R}_0|e^{\mathrm{i}H_\mathcal{R}t/\hbar}|n_\nu\rangle\langle m_\mu|e^{-\mathrm{i}H_\mathcal{R}t/\hbar}|\psi^\mathcal{R}_0\rangle, \tag{69}$$

for $t \in \mathcal{T}$. Therefore, usually, Eq. (8) can not hold if the initial vector has non-zero components in the two subspaces $\mathscr{H}_\mu$ and $\mathscr{H}_\nu$.

Then, since the time T is practically infinitely long, according to the definition of R-observable, in a practical sense, this system $\mathcal{R}$ does not have a R-observable. Without a R-observable, the assumption of MsD is not applicable to the system $\mathcal{R}$.

As an example of isolatable system, let us consider the center-of-mass (COM) degrees of freedom of a physical system, with its COM degrees of freedom taken as the system $\mathcal{R}$ and its internal degrees of freedom taken as a part of the environment of $\mathcal{R}$. If it is in principle possible for this system $\mathcal{R}$ to be uncoupled from its environment for a very long time period such that Eq. (66) holds, then, this system $\mathcal{R}$ does not have a R-observable. This implies that, in principle, quantum interference effect may be observed for the motion of the COM of appropriately prepared systems, regardless of their masses. Indeed, up to now, no upper bound has been observed experimentally for the size of a system whose COM motion may exhibit quantum interference effects.

It is straightforward to generalize the above discussions to the case that the time T is long, but not practically infinitely long. In this case, the system $\mathcal{R}$ may have a R-observable, but with a quite long decoherence time τ_d.

Finally, we give a brief discussion in some implications of the above results. The above discussions show that for a system $\mathcal{R}$ to have a R-observable, it must be non-isolatable from its environment. In other words, there should exist a part of the environment of $\mathcal{R}$, which is always accompanying $\mathcal{R}$ with sufficient interaction and is always inducing decoherence to the system $\mathcal{R}$. Some properties that the accompanying environment should have in order to guarantee the existence of a R-observable and the validity of Eq.(38), have been discussed in Sec.VII B and Sec.VII C.

In particular, Eq.(38) requires that a difference in some steps of two paths should have its decoherence effects in all the future times. Clearly, this requirement can not be met by a general Hamiltonian. In other words, it imposes a restriction to the Hamiltonian of the total system. As an example, we may consider an atom, which is always interacting with the background electromagnetic (radiation) field. Indeed, an atom has a quite specific interaction with the radiation field, determined by QED. A difficulty met in this study is the ultraviolet divergence; that is, how to treat the renormalization scheme in the study of decoherence. This is a quite complex problem and we would leave it to future investigation.

IX. APPLICATION II — AN n-LEVEL SYSTEM

In this section, we discuss a model, in which an n-level system interacts with a chaotic environment. In this model, more explicit results may be obtained following discussions given in Sec.VII B and Sec. VII C.

A. Energy eigenstates and R-observable

We consider an n-level system with normalized energy eigenstates denoted by $|\mu\rangle$,

$$H_{\mathcal{R}}|\mu\rangle = E_\mu|\mu\rangle \quad \text{for} \quad \mu = 1, 2, \cdots, n, \tag{70}$$

and projection operators $\mathcal{P}_\mu = |\mu\rangle\langle\mu|$. To find out a condition under which the corresponding observable $A_{\{\mu\}}$ can be a R-observable defined in Sec. IV B 2, we should consider a time interval $\mathcal{T} = [0, T]$, within which the non-transition condition (11) is satisfied for an initial vector of a product form, $|\Psi(0)\rangle = \left(\sum_\mu c_\mu|\mu\rangle\right)|\phi(0)\rangle$, where $|\phi(0)\rangle$ is a normalized vector in the Hilbert space of the environment $\mathcal{E}$.

In this model, off-diagonal elements $\langle\mu|\rho^{re}_{\mathcal{R}}|\nu\rangle$ with $\mu \neq \nu$ can be expressed in terms of the Loschmidt echo. To show this point, let us write Schrödinger evolution $|\Psi(t)\rangle$ in the following form,

$$|\Psi(t)\rangle = \sum_\mu c_\mu|\mu\rangle|\phi_\mu(t)\rangle. \tag{71}$$

Substituting Eq. (71) into the definition of the reduced density matrix $\rho^{re}_{\mathcal{R}}$, it is ready to find that

$$\langle\mu|\rho^{re}_{\mathcal{R}}(t)|\nu\rangle = c_\mu c^*_\nu f_{\nu\mu}(t), \tag{72}$$

where $f_{\nu\mu}(t) = \langle\phi_\nu(t)|\phi_\mu(t)\rangle$. To find out an explicit expression for $f_{\nu\mu}(t)$, one may substitute Eq. (71) into Eq. (44), getting

$$i\hbar\frac{\partial}{\partial t}|\phi_\mu(t)\rangle \doteq H^{\mathcal{E}}_\mu|\phi_\mu(t)\rangle, \tag{73}$$

where

$$H^{\mathcal{E}}_\mu = \langle\mu|H|\mu\rangle = E_\mu + H_{\mathcal{E}} + H^{\mathcal{E}}_{I\mu} \tag{74}$$

is a Hermitian operator in the Hilbert space of the environment $\mathcal{E}$, with $H^{\mathcal{E}}_{I\mu} = \langle\mu|H_I|\mu\rangle$. Hence,

$$|\phi_\mu(t)\rangle \doteq U^{\mathcal{E}}_\mu(t, 0)|\phi(0)\rangle, \tag{75}$$

where

$$U^{\mathcal{E}}_\mu(t, 0) = \exp\{-itH^{\mathcal{E}}_\mu/\hbar\}. \tag{76}$$

Then, making use of Eq. (75), it is seen that

$$f_{\nu\mu}(t) \doteq \langle\phi(0)|e^{itH^{\mathcal{E}}_\nu/\hbar}e^{-itH^{\mathcal{E}}_\mu/\hbar}|\phi(0)\rangle \tag{77}$$

is a Loschmidt-echo amplitude defined in Eq. (58).

To get an explicit estimate to $|f_{\nu\mu}(t)|$, let us consider an environment that can be modelled by a quantum chaotic system [50]. In this case, the decaying behavior of the Loschmidt echo is separated by a perturbative border ε_p, which can be estimated by [44]

$$2\pi\varepsilon_p\overline{V^2_{nd}} \sim \sigma_v\Delta, \tag{78}$$

where $\varepsilon V = H^{\mathcal{E}}_{I\nu} - H^{\mathcal{E}}_{I\mu}$ and $\overline{V^2_{nd}}$ is the average of $|\langle n|V|n'\rangle|^2$ with $n \neq n'$. Here $|n\rangle$ denote the eigenstates of $H^{\mathcal{E}}_\mu$, Δ is the mean level spacing of $H^{\mathcal{E}}_\mu$, and σ^2_v is the variance of the diagonal elements $\langle n|V|n\rangle$. Below and above the border ε_p, typically, the Loschmidt echo has a Gaussian and an exponential decay, respectively,

$$|f_{\nu\mu}(t)| \simeq e^{-\varepsilon^2\sigma^2_v t^2/2\hbar^2}, \quad \varepsilon < \varepsilon_p, \tag{79}$$

$$|f_{\nu\mu}(t)| \sim e^{-\Gamma t/2\hbar}, \quad \varepsilon > \varepsilon_p, \tag{80}$$

where $\Gamma = 2\pi\varepsilon^2\overline{V^2_{nd}}/\Delta$ [43–46].

For a large environment, Δ is small, hence, the border ε_p is low and usually one is interested in the case of $\varepsilon > \varepsilon_p$. In this case, the Loschmidt echo has the exponential decay in Eq. (80), as a result, $\langle\mu|\rho^{re}_{\mathcal{R}}(t)|\nu\rangle$ becomes negligibly small for times beyond a decoherence time τ_d,

$$\tau_d = k\hbar\Delta/[\pi\varepsilon^2\overline{V^2_{nd}}], \tag{81}$$

where k is a number determined by the accuracy required. This exponential decay of the echo stops when its saturation value is reached, which is inversely proportional to the dimension N of the Hilbert space of the environment [46]. For a sufficiently large environment, the saturation value is (effectively) zero.

Finally, let us discuss the condition for $A_{\{\mu\}}$ to be a R-observable. First, in the case of a constant V, $|f_{\nu\mu}(t)| = 1$ [see Eq. (77)] and $\tau_d = \infty$ [see Eq. (81)], hence, there is no decoherence induced by the environment and Eq. (8) can never be satisfied; in this case, $A_{\{\mu\}}$ can not be a R-observable. Second, for a non-constant V with a non-zero $\overline{V^2_{nd}}$, the Loschmidt echo decays with time, characterized by the decoherence time τ_d given in Eq. (81); in this case, $A_{\{\mu\}}$ can be a R-observable.

Summarizing the above discussions, we reach the following conclusion: For an environment that can be modeled by a quantum chaotic system, $A_{\{\mu\}}$ is a R-observable of the system $\mathcal{R}$, if the environment is sufficiently large and there are sufficient differences among $H^{\mathcal{E}}_{I\mu}$.

B. $D_{\beta\alpha}$ expressed in the Hilbert space of the environment

The condition under which Eq.(59) may be satisfied can be discussed in a way similar to that given in Sec. VII C. In the present model, we may express $D_{\beta\alpha}$ in terms of quantities in the Hilbert space of the environment.

For the simplicity in discussion, let us consider an initial vector $|\Psi(t_0)\rangle = |\mu_0\rangle|\phi_0\rangle$, with $|\phi_0\rangle \in \mathscr{H}_{\mathcal{E}}$. Making use of Eq. (61), the component $|\Psi_\alpha(t)\rangle$ can be written as

$$|\Psi_\alpha(t)\rangle = \sum_\mu |\mu\rangle|\phi^\alpha_\mu(t)\rangle, \tag{82}$$

where

$$|\phi^\alpha_\mu(t)\rangle = \langle\mu|W_\alpha(t,t_0)|\mu_0\rangle|\phi_0\rangle. \tag{83}$$

Making use of Eq. (62), we find that

$$\langle\mu|W_\alpha(t,t_0)|\mu_0\rangle = Y_{\mu\mu^\alpha_{(n)}}(t,\widetilde{\tau}^\alpha_n))\ U^{\mathcal{E}}_{\mu^\alpha_{(n)}}(\widetilde{\tau}^\alpha_n,\tau^\alpha_n)\cdots$$
$$\cdot U^{\mathcal{E}}_{\mu^\alpha_{(2)}}(\widetilde{\tau}^\alpha_2,\tau^\alpha_2)Y_{\mu^\alpha_{(2)}\mu^\alpha_{(1)}}(\tau^\alpha_2,\widetilde{\tau}^\alpha_1)U^{\mathcal{E}}_{\mu^\alpha_{(1)}}(\widetilde{\tau}^\alpha_1,\tau^\alpha_1)Y_{\mu^\alpha_{(1)}\mu_0}(\tau^\alpha_1,t_0), \tag{84}$$

where $U^{\mathcal{E}}_\mu(t,t')$ is defined like the one in Eq. (76) and

$$Y_{\mu\mu'}(t,t') = \langle\mu|U(t,t')|\mu'\rangle. \tag{85}$$

The operator $Y_{\mu\mu'}$, though an operator in $\mathscr{H}_{\mathcal{E}}$, in fact represents transition between subspaces $\mathscr{H}_\mu$ and $\mathscr{H}_{\mu'}$ in the total Hilbert space.

Thus, $D_{\beta\alpha} = \langle\Psi_\beta(t)|\Psi_\alpha(t)\rangle$ has the following expression in the Hilbert space of the environment,

$$D_{\beta\alpha} = \sum_\mu \langle\phi_0|Y^\dagger_{\mu^\beta_{(1)}\mu_0} U^{\mathcal{E}\dagger}_{\mu^\beta_{(1)}} Y^\dagger_{\mu^\beta_{(2)}\mu^\beta_{(1)}} U^{\mathcal{E}\dagger}_{\mu^\beta_{(2)}} \cdots U^{\mathcal{E}\dagger}_{\mu^\beta_{(m)}} Y^\dagger_{\mu\mu^\beta_{(m)}}$$
$$\cdot Y_{\mu\mu^\alpha_{(n)}} U^{\mathcal{E}}_{\mu^\alpha_{(n)}} \cdots U^{\mathcal{E}}_{\mu^\alpha_{(2)}} Y_{\mu^\alpha_{(2)}\mu^\alpha_{(1)}} U^{\mathcal{E}}_{\mu^\alpha_{(1)}} Y_{\mu^\alpha_{(1)}\mu_0}|\phi_0\rangle, \tag{86}$$

where the dependence of $U^{\mathcal{E}}_\mu$ and $Y_{\mu\mu'}$ on times is not written explicitly.

X. APPLICATION III — IRREVERSIBILITY OF BRANCHING PROCESSES

In this section, we discuss the effectively-irreversible feature of the proposed theory and, as an application, we use the theory to derive a master equation for an appropriate process.

A. Irreversible feature of time evolution

The branching picture of time evolution discussed in Sec.V A is not time-reversible. This is not in confliction with the time-reversal symmetry of Schrödinger equation, because the branching picture represents a property of physical states only for initial conditions satisfying the Initial-vector-restriction.

A quantity that can characterize the irreversible feature of branching is von Neumann entropy for the total system,

$$S(t) = -\mathrm{Tr}\{\rho\ln\rho\}. \tag{87}$$

Substituting the mixed-state description $\rho_\Upsilon(t)$ in Eq. (21) into Eq. (87) and making use of Eq. (22) for the probability for the realization of a path α, we have

$$S_\Upsilon(t) = -\sum_\alpha P_\alpha(t)\ln P_\alpha(t). \tag{88}$$

When no branch-splitting happens, $S_\Upsilon(t)$ keeps constant as a result of Schrödinger evolution; while, at each splitting time t^α_i along a path $\alpha \in \Upsilon$, the entropy $S_\Upsilon(t)$ obtains a discontinuous increment. Therefore, $S_\Upsilon(t)$ may increase but never decrease with increasing time. Interestingly, since the system $\mathcal{R}$ may get some new definite value μ at each splitting point, each increment of the von Neumann entropy is accompanied with the possibility of a measurement, i.e., the possibility of gathering information.

One may note the following differences of the entropy $S_\Upsilon(t)$ in Eq. (88) from the thermodynamic entropy. That is, $S_\Upsilon(t)$ increases without any upper bound; it is usually un-measurable, since different branches may have the same value of μ. An entropy that may have an upper bound will be discussed in the next section.

B. Master equation for an ideal case of branching

In this section, as an application of the theory proposed above, we derive a master equation for the probabilities for the system $\mathcal{R}$ to take definite values of μ.

For the simplicity in discussion, we consider an ideal case, in which all the time intervals $[\tau^\alpha_{i+1},\widetilde{\tau}^\alpha_i]$ are very short such that the non-transition condition is satisfied for almost all times along the paths. This is a good approximation in some practical situations. For the same reason, we assume that the branching times t^α_i, as well as the times τ^α_i and $\widetilde{\tau}^\alpha_i$ are path-independent, thus, we can drop the superscript α in the labeling of these times. Moreover, we consider one R-observable $A_{\{\mu\}}$ only.

Since the non-transition condition is satisfied for almost all the times, the system $\mathcal{R}$ has a definite value of μ for almost all the times along the paths. We use $p_\mu(t)$ to denote the probability for $\mathcal{R}$ to have a definite value μ at a time t,

$$p_\mu(t) = \sum_{\alpha \text{ with } |\Psi_\alpha(t)\rangle\in\mathscr{H}_\mu} P_\alpha(t). \tag{89}$$

Using this quantity, we can define an entropy for the measuring apparatus $\mathcal{R}$,

$$S_{\mathcal{R}}(t) = -\sum_\mu p_\mu(t)\ln p_\mu(t). \tag{90}$$

This entropy may have an upper bound and is in principle measurable.

Below, we derive a master equation for the time evolution of $p_\mu(t)$. Let us consider a path α that ends at a time t beyond a branching time t_n, with $|\Psi_\alpha(t)\rangle$ expressed in Eq. (20). At a time t_{n+1}, this path splits, resulting in paths we denote by $|\Psi_\beta(t)\rangle$ for t beyond t_{n+1}, with

$$\beta = (\alpha \to \mu_{(n+1)}(t_{n+1})), \tag{91}$$

$$|\Psi_\beta(t)\rangle = U(t, t_{n+1})\mathcal{P}_{\mu_{(n+1)}} U(t_{n+1}, t_n)|\Psi_\alpha(t_n)\rangle. \tag{92}$$

The probability for the realization of a path β, $P_\beta(t_{n+1}) = \langle\Psi_\beta(t)|\Psi_\beta(t)\rangle$, can be written as

$$P_\beta(t_{n+1}) = \Gamma_n(\alpha, \mu_{(n+1)})P_\alpha(t_n), \tag{93}$$

where $\Gamma_n(\alpha, \mu_{(n+1)})$ is defined by

$$\Gamma_n(\alpha, \mu) = \frac{1}{P_\alpha(t_n)}\langle\Psi_\alpha(t_n)|U^\dagger(t_{n+1}, t_n)\mathcal{P}_\mu U(t_{n+1}, t_n)|\Psi_\alpha(t_n)\rangle. \tag{94}$$

From Eq. (89) we have

$$p_\mu(t_n) = \sum_{\alpha \text{ with } \mu^\alpha_{(n)}=\mu} P_\alpha(t_n). \tag{95}$$

For a given value of $\mu_{(n+1)}$, say, μ, the path β is determined by the path α, hence, similar to Eq. (95), for the time t_{n+1} we have

$$p_\mu(t_{n+1}) = \sum_\alpha P_\beta(t_{n+1})\big|_{\mu_{(n+1)}=\mu}. \tag{96}$$

We denote by $\Gamma_n(\mu', \mu)$ the average of $\Gamma_n(\alpha, \mu)$ over those paths α that have a given value μ' of $\mu^\alpha_{(n)}$,

$$\Gamma_n(\mu', \mu) = \frac{1}{\mathcal{N}} \sum_{\alpha \text{ with } \mu^\alpha_{(n)}=\mu'} \Gamma_n(\alpha, \mu), \tag{97}$$

where

$$\mathcal{N} = \sum_{\alpha \text{ with } \mu^\alpha_{(n)}=\mu'} 1. \tag{98}$$

Then, we can write

$$\Gamma_n(\alpha, \mu) = \Gamma_n(\mu^\alpha_{(n)}, \mu) + \delta\Gamma_n(\alpha, \mu), \tag{99}$$

where $\delta\Gamma_n(\alpha, \mu)$ denotes deviation of $\Gamma_n(\alpha, \mu)$ from its average value $\Gamma_n(\mu^\alpha_{(n)}, \mu)$, with average taken over paths α having the same value of $\mu^\alpha_{(n)}$. Substituting Eq. (99) with $\mu = \mu_{(n+1)}$ into Eq. (93), then, substituting the result into Eq. (96), we have

$$p_\mu(t_{n+1}) = \sum_\alpha \Gamma_n(\mu^\alpha_{(n)}, \mu)P_\alpha(t_n) + \Delta p, \tag{100}$$

where

$$\Delta p = \sum_\alpha \delta\Gamma_n(\alpha, \mu)P_\alpha(t_n). \tag{101}$$

The summation over all the paths α is equivalent to a summation over those paths α with a fixed value of $\mu^\alpha_{(n)} = \mu'$, followed by a summation over μ'. Hence, making use of Eq. (95), from Eq. (100) we have

$$p_\mu(t_{n+1}) = \sum_{\mu'} \Gamma_n(\mu', \mu)p_{\mu'}(t_n) + \Delta p. \tag{102}$$

To give an estimate to Δp, we note that by definition the average of $\delta\Gamma_n(\alpha, \mu)$ is zero and $\sum_\alpha P_\alpha(t) = 1$. This implies that, when the number of α is sufficiently large, Δp is usually negligibly small. In fact, in the case that $\delta\Gamma_n(\alpha, \mu)P_\alpha(t_n^+)$ can be regarded as a random number, one has $\Delta p \sim 1/\sqrt{M_n}$, where M_n is the number of the paths α. It is easy to see that M_n increases exponentially with increasing n.

To summarize, when n, the number of steps is sufficiently large, the probability for $\mathcal{R}$ to take a definite value of μ at t_{n+1} satisfies a master equation,

$$p_\mu(t_{n+1}) \simeq \sum_{\mu'} \Gamma_n(\mu', \mu)p_{\mu'}(t_n). \tag{103}$$

From the definitions given in Eqs. (94) and (97), it is easy to check that $\Gamma_n(\mu', \mu) \geq 0$ and

$$\sum_\mu \Gamma_n(\mu', \mu) = 1. \tag{104}$$

Some remarks: Compared with derivations of master equations given in the usual quantum theory, the derivation given above has the following advantages: It is relatively simple and uses less approximations (we do not need to use approximations like Born approximation and Markov approximation).

XI. DISCUSSIONS AND CONCLUSIONS

In this section, we first give comparisons of the proposed theory with CHI and MWI of quantum mechanics. Then, we give a brief summary for the main results of this paper, as well as some discussions.

A. Comparison with CHI of quantum mechanics

In this section, we discuss relations between the theory proposed here and CHI. In CHI, the time evolution of a quantum system has a stochastic nature and is described by (quantum) consistent histories [8–16]. Each history is composed of a sequence of events represented by time-ordered projection operators, with unitary connection between each two successive events. The consistency of consistent histories is determined by a consistency condition. One projective decomposition of the identity operator is called a framework. A single framework rule must be obeyed when CHI is used, which states that a valid description must use one framework only, even though other frameworks are also legitimate.

One may note some similarities between the mathematical formulations of some main results of the theory proposed here and of CHI, which we list below.

(1) A description given by a path in the theory here, namely, $|\Psi_\alpha(t)\rangle$ in Eq. (20), has a formal similarity to the contribution given by a history in CHI.

(2) Substituting Eq. (20) into Eq. (60), it is seen that the quantity $\mathcal{D}_{\alpha\alpha'}$ can be written in a form with formal similarity to the so-called decoherence functional $\mathcal{D}(\beta,\beta')$ in CHI [14, 15],

$$\mathcal{D}(\beta,\beta') = \mathrm{Tr}\Big[P^{(n)}_{\beta_n}U(t_n,t_{n-1})\dots P^{(1)}_{\beta_1}U(t_1,t_0)\rho(t_0) \\ U^\dagger(t_1,t_0)P^{(1)}_{\beta'_1}\cdots U^\dagger(t_n,t_{n-1})P^{(n)}_{\beta'_n}\Big], \quad (105)$$

where β indicates a history and $P^{(j)}_{\beta_j}$ of $j=1,\dots,n$ denote projection operators in the history β. The two quantities $\mathcal{D}_{\alpha\alpha}$ and $\mathcal{D}(\beta,\beta)$ give the corresponding probabilities, respectively, in the two theories.

(3) One of the two requirements of the principle of compatible description, namely, Eq. (38) has the same formal form as the consistency condition in CHI, which is

$$\mathcal{D}(\beta,\beta') = \delta_{\beta\beta'}\mathcal{D}(\beta,\beta). \quad (106)$$

However, despite the formal similarities mentioned above, the two theories have profound differences in their physical contents, as listed below.

(i) *Most of the consistent-histories descriptions allowed in CHI do not have any corresponding description in the theory proposed here.* The reason is as follows. In CHI, to have a consistent-histories description of the total system, the only restriction is given by the consistency condition in Eq.(106) related to some instants (not necessarily for all the times). It has been found that many of the consistent-histories descriptions do not have a quasiclassical feature [51] (see also discussions given in Ref.[9]). While, the theory proposed here has more stringent restrictions to descriptions of physical states of the total system: (a) only R-observables can be used in the assumption of MsD to get a mixed-state description, (b) the non-transition condition must be satisfied around each branching point, and (c) the principle of compatible description must be obeyed for all the times.

For example, in CHI, one is allowed to use a few projection operators related to a few instants to construct consistent histories, as long as the histories satisfy the consistency condition in Eq.(106) for these instants, in spite of what may happen in the future times. However, usually, such obtained consistent-histories descriptions do not have corresponding descriptions in the theory proposed here, because (i) the projection operators considered in CHI are not required to correspond to a R-observable in the theory here, (ii) the non-transition condition is not required to be satisfied around the instants considered in CHI, (iii) more importantly, the descriptions allowed in CHI do not necessarily satisfy the principle of compatible description, i.e., Eqs.(38) and (30), in all the future times.

(ii) *There is no single-framework rule in the theory proposed here*, while, the single-framework rule must be obeyed in CHI to avoid logical inconsistency. (There have been some debates in its physical validity [30, 31].) In fact, in the theory here, the system $\mathcal{R}$ has a finest R-observable, with other R-observables of $\mathcal{R}$ being its coarse-grainings; in the language of CHI, this implies that there exists only one legitimate (finest) 'framework' for the system $\mathcal{R}$.

(iii) The Initial-vector-restriction may effectively break the time-reversal symmetry of Schrödinger equation; while, the time-reversal symmetry is maintained in CHI.

(iv) In CHI, the consistency condition in Eq.(106) is introduced to guarantee the validity of the sum rule of probability. In the theory here, the principle of compatible description states the physical compatibility of different mathematical descriptions for the same physical state of the total system.

B. Comparison with MWI of quantum mechanics

The MWI of quantum mechanics has two main assumptions [5–7]. Namely, (i) Schrödinger equation holds universally, and (ii) the state vector of the total system splits constantly into branches. One may combine the MWI and the decoherence theory to get a more complete picture for the time evolution, with a branch in MWI related to a preferred (pointer) state (or subspace) in the decoherence theory (see, e.g., Ref.[28]).

There also exist some formal similarities between the theory proposed here and the combination of MWI and decoherence theory.

(1) Schrödinger equation gives the dynamical law in both theories.

(2) By virtue of the assumption of MsD (Part II of the third assumption), the theory here gives a branching picture of time evolution, as illustrated in Fig.2, which has a formal similarity to that in MWI.

(3) The concept of R-observable has a close relationship to the concept of preferred basis (subspace) in the decoherence theory [2, 4, 27, 28, 48, 49, 52–57], though not exactly the same (see Sec.B of the appendix for further discussions).

Meanwhile, the two theories have the following main differences:

(i) In the theory here, the non-transition condition gives an explicitly expressed condition for branching to happen. In MWI, there is no such a condition.

(ii) MWI does not have a counterpart of Part I of the third assumption here, namely, the principle of compatible description.

(iv) In the theory here, there exists only one real world; the description of the world may split into branches, but, the real world never splits. In MWI, the world may split into many worlds.

Discussions given in this section and the previous section show that the theory proposed here can be regarded as certain type of unification of CHI and MWI+decoherence theory [58]. However, it is not a direct unification, since the theory here abandons both the single framework rule in CHI and the assumption about the splitting of the real world in MWI. In addition, the theory here proposes a concrete condition (non-transition condition) for the appearance of definite properties, which is given in neither CHI nor MWI.

C. Summary and discussions

In this paper, we have proposed a quantum theory for a total system including one internal measuring apparatus. The theory is based on three basic assumptions, which roughly speaking have the following contents: (i) the Hilbert space as the state space, (ii) Schrödinger equation as the dynamical law, and (iii) the assumption of MsD for vectors satisfying the Initial-vector-restriction. The assumption of MsD states that some pure-vector descriptions of the total system may imply the existence of certain mixed-state descriptions. The Initial-vector-restriction is a mathematical expression of the principle of compatible description, which states that different mathematical descriptions for the same physical state of the total system must give consistent predictions for measurement results of the internal measuring apparatus.

Loosely speaking, the proposed theory gives the following descriptions for the total system. Starting from an initial vector, there always exists a pure vector description $|\Psi(t)\rangle$, given by Schrödinger equation. Usually, this pure vector description does not directly give predictions for definite properties of the internal measuring apparatus. For a physical state of the total system that is initially described by a vector satisfying the Initial-vector-restriction, the pure vector description $|\Psi(t)\rangle$ may be physically equivalent to certain mixed-state descriptions at some times. Such a mixed-state description may predict some definite property of the internal measuring apparatus.

The above discussed Initial-vector-restriction imposes a restriction to initial vectors in the Hilbert space. It may effectively break the time-reversal symmetry of Schrödinger equation, leading to the irreversibility of some processes, since the time reversal of a vector satisfying the Initial-vector-restriction does not necessarily satisfy the Initial-vector-restriction. This result may shed new light in the old problem of the microscopic origin of the macroscopic irreversibility stated in the second law of thermodynamics.

One characteristic feature of the theory proposed in this paper, distinguishing it from other theories, is the role played by the non-transition condition in determining the existence of a definite property of the internal measuring apparatus. Due to this feature, it is possible that the proposed theory might give some experimentally testable predictions, which are not equivalent to the corresponding predictions of other quantum theories. However, lots of work are needed before a concrete scheme for experimental test can be constructed for this purpose, in particular, because of the difficulty of finding solutions of Eq.(38).

In the case that a physical state of a system may have more than one mathematical descriptions, the physical equivalency of the descriptions may have some non-trivial implications in properties of the system. For example, The Lorentz invariance in the theory of relativity and the gauge symmetry in the quantum field theory have quite rich implications. In the theory proposed here, the physical equivalency of some pure-vector and mixed-state descriptions requires Eq.(38). A natural and interesting question is whether there may exist other non-trivial consequence(s) of this equivalency, which deserves investigation in the future.

Acknowledgments

The author is grateful to A.J. Leggett, Yan Gu, Hong Zhao, C. Kiefer, and Jiangbin Gong for valuable discussions and suggestions. This work is partially supported by Natural Science Foundation of China under Nos. 10775123 and 10975123.

Appendix A: A relation between trees and their coarse-grainings

In this appendix, we show that, to obtain a valid coarse-grained tree, in addition to the coarse-graining at one step of a path of the original tree, modifications in the subsequent steps may also be needed.

It would be convenient to write explicit dependence of a tree on the initial vector $|\Psi(t_0)\rangle$ and on the ending time t, i.e., write it as $\Upsilon(|\Psi(t_0)\rangle, t)$. Let us consider an intermediate time $\widetilde{t_0}$, at which the tree Υ gives components $|\Psi_\alpha(\widetilde{t_0})\rangle$. Taking a component $|\Psi_\alpha(\widetilde{t_0})\rangle$ as an initial condition, the subsequent evolution in the tree $\Upsilon(|\Psi(t_0)\rangle, t)$ forms a sub-tree, denoted by $\Upsilon(|\Psi_\alpha(\widetilde{t_0})\rangle, t)$. This possibility can be seen clearly in Fig. 2. The relation between the original tree and the sub-trees can be written as

$$\Upsilon(|\Psi(t_0)\rangle, t) = \bigcup_{\alpha(\widetilde{t_0})} \alpha(\widetilde{t_0}) \odot \Upsilon(|\Psi_\alpha(\widetilde{t_0})\rangle, t), \tag{A1}$$

where we write explicitly the ending time of the paths α and use $\odot$ to indicate a successive relationship of a path and a following sub-tree.

Let us use B to denote the next splitting point of a path $\alpha(\widetilde{t_0})$, which takes place at a time t_B and is related with a R-observable $A_{\{\mu\}}$; we use $\alpha_B(t)$ of $t \in [\widetilde{t_0}, t_B]$ to indicate the extension of the path $\alpha(\widetilde{t_0})$. For B to be a splitting point, the non-transition condition should be satisfied for the R-observable $A_{\{\mu\}}$ around the time t_B. It is easy to see that, around t_B, the non-transition condition is also satisfied for an arbitrary coarse-grained R-observable $A_{\{\eta\}}$ defined using Eq. (9). If at the point B we use $A_{\{\eta\}}$, instead of $A_{\{\mu\}}$, to generate the splitting of the component $|\Psi_{\alpha_B}(t_B)\rangle$, we will get a coarse-grained sub-tree, denoted by $\Upsilon_c(|\Psi_{\alpha_B}(\widetilde{t_0})\rangle, t)$. Then, we get the following coarse-grained tree Υ_c for the initial condition $|\Psi(t_0)\rangle$,

$$\begin{aligned} \Upsilon_c(|\Psi(t_0)\rangle, t) = {} & \alpha_B(\widetilde{t_0}) \odot \Upsilon_c(|\Psi_{\alpha_B}(\widetilde{t_0})\rangle, t) \\ & \bigcup_{\alpha \neq \alpha_B} \alpha(\widetilde{t_0}) \odot \Upsilon(|\Psi_\alpha(\widetilde{t_0})\rangle, t), \end{aligned} \tag{A2}$$

where terms in the second line are the same as the corresponding ones in Eq. (A1).

Let us compare the sub-tree $\Upsilon(|\Psi_{\alpha_B}(\widetilde{t_0})\rangle, t)$ and the coarse-grained sub-tree $\Upsilon_c(|\Psi_{\alpha_B}(\widetilde{t_0})\rangle, t)$. They have the same initial condition $|\Psi_{\alpha_B}(\widetilde{t_0})\rangle$ and the same first splitting time t_B, but they have different R-observables at the splitting time t_B, namely, $A_{\{\mu\}}$ and $A_{\{\eta\}}$, respectively. Since $A_{\{\eta\}}$ is a coarse-graining of $A_{\{\mu\}}$, the components of the two sub-trees at a time t immediately beyond the splitting time t_B have the following relation,

$$|\Psi_{\widetilde{\alpha}}(t)\rangle = \mathcal{P}_\mu |\Psi_{\widetilde{\alpha}_c}(t)\rangle, \tag{A3}$$

where we use $\widetilde{\alpha}$ and $\widetilde{\alpha}_c$ to denote paths in the two sub-trees, respectively. Usually, $|\Psi_{\widetilde{\alpha}}(t)\rangle$ is not equal to $|\Psi_{\widetilde{\alpha}_c}(t)\rangle$, hence, if $|\Psi_{\widetilde{\alpha}}(t)\rangle$ satisfies the non-transition condition at a time $\widetilde{t_2}$ for some R-observable, it is not necessary for $|\Psi_{\widetilde{\alpha}_c}(t)\rangle$ to satisfy the non-transition condition at the same time and for the same R-observable. As a result, the second splitting points of the two sub-trees may be different. Therefore, a mere replacement of $\mu(t_B)$ by $\eta(t_B)$ at the point B, without any change in the following steps, does not necessarily give the coarse-grained tree $\Upsilon_c(|\Psi_{\alpha_B}(\widetilde{t_0})\rangle, t)$.

Appendix B: R-observable and preferred pointer basis

The basic physical idea behind the concept of R-observable is similar to that behind the concept of preferred (pointer) basis in the decoherence theory [2, 28, 48, 49, 52, 53], more exactly, to its generalization as preferred subspace [54]. The two concepts have the following main difference: The decoherence property of a R-observable given in Eq.(8) is required to have certain initial-condition independence and to hold when the non-transition condition is satisfied. In the decoherence theory, although some type of initial-condition independence is required in some studies of preferred basis, it is not a general requirement; the non-transition condition is not a general requirement, either.

To be more specific, in the decoherence theory, in the weak coupling limit of the system-environment interaction, energy eigenstates of a system with discrete energy levels form a preferred basis when some condition is satisfied [49, 53]. While, for a quantum Brownian particle with weak coupling with the environment, coherent states have been found to be preferred states [4, 28, 55, 56]. Furthermore, in the strong coupling limit, eigenstates of the system-environment interaction Hamiltonian may form a preferred basis [27, 53, 57].

Since the non-transition condition generally implies Eq.(49), the weak coupling limit of the system-environment interaction is a special case of what we are considering here. Indeed, as shown in Sec.IX, the finest R-observable of a system $\mathcal{R}$ with n discrete energy levels may be related to the energy eigenstates, similar to the case of preferred basis in the weak coupling limit in the decoherence theory. However, as shown in Sec.VIII, the COM degrees of freedom of a quantum Brownian particle does not have a R-observable, in contrast to the case in the decoherence theory with coherent states as preferred states.

[1] J. von Neumann, *Mathematische Grundlagen der Quantummechanik*, (Springer, Berlin, 1932). English translation: *Mathematical Foundations of Quantum Mechanics*, (Princeton University press, Princeton, NJ, 1955).

[2] M. Schlosshauer, Rev. Mod. Phys. **76**, 1267 (2004).

[3] F. Laloë, Am. J. Phys. **69**, 655 (2001).

[4] E. Joos, H.D. Zeh, C. Kiefer, D. Giulini, J. Kupsch, and I.-O. Stamatescu, *Decoherence and the Appearance of a Classical World in Quantum Theory*, 2nd ed., (Springer, Berlin, 2003).

[5] H. Everett, Rev. Mod. Phys. **29**, 454 (1957).

[6] *The Many-Worlds Interpretation of Quantum Mechanics*, edited by B.S. DeWitt and N. Graham, (Princeton University Press, Princeton, 1973).

[7] D. Deutsch, Int. J. Theor. Phys. **24**, 1 (1985).

[8] R. B. Griffiths, J. Stat. Phys. **36**, 219 (1984).

[9] R. B. Griffiths, Phys. Rev. A **54**, 2759 (1996); *ibid.* **57**, 1604 (1998).

[10] R.G. Griffiths, *Consistent Quantum Theory* (Cambridge University Press, Cambridge, United Kingdom, 2002).

[11] R. Omnès, J. Stat. Phys. **53**, 893 (1988).

[12] R. Omnès, Rev. Mod. Phys. **64**, 339 (1992).

[13] R. Omnès, *Understanding Quantum Mechanics* (Princeton University Press, Princeton, New Jersey, 1999).

[14] M. Gell-Mann and J. Hartle, in *Proceedings of the 3rd International Symposium on the Foundations of Quantum Mechanics in the Light of New Technology*, Physical Society of Japan, Tokyo,1990, S. Kobayashi et al. Eds.; in *Complexity, Entropy, and Physics of Information*, W. Zurek ed. (Addison Wesley, Reading, 1990).

[15] M. Gell-Mann and J. Hartle, Phys. Rev. D **47**, 3345 (1993).

[16] P. Hohenberg, arXiv:0909.2359.

[17] L. de Broglie, *An Introduction to the Study of Wave Mechanics*, (E. Physica Dutton and Co., New York, 1930).

[18] D. Bohm, Phys. Rev. **85**, 166 (1952); **85**, 180 (1952).

[19] D. Bohm and J. Bub, Rev. Mod. Phys. **38**, 453 (1966).

[20] G.C. Ghirardi, A. Rimini, and T. Weber, in *Quantum Probabililty and Applications II*, Vol 1136 of *Lecture Notes in Mathematics*, edited by L. Accardi and W. von Waldenfels (Springer, Berlin, 1985).

[21] G.C. Ghirardi, A. Rimini, and T. Weber, Phys. Rev. D **34**, 470 (1986).

[22] P. Pearle, Phys. Rev. A **39**, 2277 (1989).

[23] G.C. Ghirardi, P. Pearle, and A. Rimini, Phys. Rev. A **42**, 78 (1990).

[24] S.L. Adler and A. Bassi, Science **325**, 275 (2009).

[25] A. Bassi and G.C. Ghirardi, Phys. Rep. **379**, 257 (2003).

[26] H.D. Zeh, Found. Phys. **1**, 69 (1970); *ibid.* **3**, 109 (1973).

[27] W. H. Zurek, Phys. Rev. D **24**, 1516 (1981); *ibid.* **26**, 1862 (1982).

[28] W.H. Zurek, Rev. Mod. Phys. **75**, 715 (2003).

[29] A.J. Leggett, J.Phys. Cond.Mat., **14**, R415 (2002); Prog. Theor. Phys. supplement, **170**, 100 (2007).

[30] A. Bassi and G.C. Ghirardi, Phys. Lett. A **257**, 247 (1999); Phys. Lett. A **265**, 153 (2000); J. Stat. Phys. **98**, 457 (2000); J. Stat. Phys. **99**, 1427 (2000).

[31] R.B. Griffiths, Phys. Lett. A **265**, 12 (2000); J. Stat. Phys. **99**, 1409 (2000).

[32] A.Kent and J.McElwaine, Phys.Rev.A **55**, 1703 (1997).

[33] In the standard formalism of quantum mechanics, in order to explain experimental results, the axiom of measurement is introduced, which states that a measurement on an observable of a system leads the state of the system to an eigenstate of the observable (collapse of state vector). As pointed out by many authors, this axiom of measurement can not be directly generalized to the case of a total system possessing an internal measuring apparatus.

[34] Relation between descriptions of a total system given by two independent internal observers is a difficult topic and we would leave it to future investigation. In fact, to know the relation, communication between the two observers is necessary. Since a measurement should be carried out for a successful communication, communications may change the state of the total system. This makes the case of two observers much more complex than the case of one observer considered here and we would leave this topic for future investigation.

[35] In the usual quantum mechanics for an external observer, the difference between a mixed-state description and a pure-vector description for the same system is in principle experimentally testable by an external measuring apparatus, therefore, the two descriptions can not describe the same physical state of the system.

[36] When assigning definite properties to a subsystem of a total system under Schrödinger evolution, it is usually unavoidable to face the problem of small probability. For example, in the decoherence approach to the foundations of quantum mechanics [2], decoherence may reduce off-diagonal elements of a reduced density matrix to small quantities, but can hardly make them exact zero. The theory proposed here belongs to this type of theory.

[37] Obviously, consideration of initial vectors of the product form is not the only possible choice. But, for the purpose here, i.e., to determine the projection operators $\mathcal{P}_\mu$ (R-observable), the consideration of these vectors may be enough.

[38] I. Prigogine and I. Stengers, *Order Out of Chaos*, (Bantam Books Inc., 1984).

[39] H. D. Zeh, *The Physical Basis of the Direction of Time*, 3th ed. (Springer, Berlin, 1999).

[40] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* , (Oxford University Press, New York, 2002).

[41] A. Peres, Phys. Rev. A **30**, 1610 (1984).

[42] T. Gorin, T. Prosen, T.H. Seligman, and M. Žnidarič, Phys. Rep. **435**, 33 (2006) and references there in.

[43] R.A. Jalabert and H.M. Pastawski, Phys. Rev. Lett. **86**, 2490 (2001); Ph. Jacquod, P.G. Silvestrov, and C.W.J. Beenakker, Phys. Rev. E **64**, 055203(R) (2001); F.M. Cucchietti, *et al*, Phys. Rev. E **65**, 046209 (2002).

[44] N. R. Cerruti and S. Tomsovic, J. Phys. A **36**, 3451 (2003); Phys. Rev. Lett. **88**, 054103 (2002).

[45] W.-G Wang and B. Li, Phys. Rev. E **66**, 056208 (2002); W.-G. Wang, G. Casati, and B. Li, *ibid.* **69**, 025201(R)(2004); W.-G. Wang, *et al*, *ibid.* **71**, 037202 (2005); W.-G Wang and B. Li, *ibid.* **71**, 066203 (2005).

[46] T. Prosen and M. Žnidarič, J. Phys. A **35**, 1455 (2002); G. Benenti and G. Casati, Phys. Rev. E **65**, 066205(2002).

[47] Wen-ge Wang, Pinquan Qin, Lewei He, and Ping Wang, Phys. Rev. E, **81**, 016214 (2010).

[48] T. Gorin, T. Prosen, T.H. Seligman, and W.T. Strunz, Phys. Rev. A **70**, 042105 (2004).

[49] Wen-ge Wang, Jiangbin Gong, G. Casati, and Baowen Li, Phys. Rev. A **77**, 012108 (2008).

[50] M. Srednicki, Phys. Rev. E **50**, 888 (1994).

[51] F. Dowker and A. Kent, Phys. Rev. Lett. **75**, 3038 (1995); J. Stat. Phys. **82**, 1575 (1996).

[52] D. Spehner and F. Haake, J. Phys. A **41**, 072002 (2008); Phys. Rev. A **77**, 052114 (2008).

[53] J. P. Paz and W. H. Zurek, Phys. Rev. Lett. **82**, 5181 (1999).

[54] A. Peres, Phys. Rev. A **61**, 022116 (2000).

[55] W. H. Zurek, S. Habib, and J. P. Paz, Phys. Rev. Lett. **70**, 1187 (1993).

[56] J. Eisert, Phys. Rev. Lett. **92**, 210401 (2004); H. M. Wiseman and J. A. Vaccaro, Phys. Rev. A **65**, 043606 (2002).

[57] D. Braun, F. Haake, and W.T. Strunz, Phys. Rev. Lett. **86**, 2913 (2001).

[58] The possibility of making use of some results of the decoherence theory in the CHI approach has been discussed

by J. P. Paz and W. H. Zurek in Phys. Rev. D **48**, 2728 (1993). It has been shown that consistent histories may become more stable, if the time separation of events is required to be larger than the typical decoherence time and the projection operators for events are required to be associated with preferred basis.